

Exam FY3106 / FY8303

December 2021

Problem 1

a) H is dimensionless: $Z = \int D\theta e^{-H}$

$$\text{Thus: } [g] \cdot L^{d-2} (L^0)^2 = L^0$$

$$[g] = L^{2-d}$$

Scaling-dimension of g : $2-d$

b) $| \langle \psi \rangle |^2 \sim 1/L^{2\beta}$ ($| \langle \psi \rangle | \sim 1/L^\beta$)

One way of arriving at H would be to start with

$$E = \frac{\hbar^2}{2m} |\nabla \psi|^2 + \alpha |\psi|^2 + \frac{u}{2} |\psi|^4$$

with $\psi = |\psi| e^{i\theta} \approx |\psi_0| e^{i\theta}$

$$\text{Then } g = \frac{\hbar^2}{m} |\psi_0|^2$$

$$g \sim |\psi_0|^2 \text{ density}$$
$$| \langle \psi \rangle |^2 \sim \text{density}$$

$$\gamma \sim g \quad (\text{unnormalised})$$

Thus, if $\gamma \sim |\psi|^\omega$

then it is natural to compare

$$\omega \text{ to } 2\beta$$

since $g \sim |\psi|^2$

$$c) \quad \gamma \sim g \sim \int |\psi|^{2-d} \sim |\psi|^{\nu(d-2)}$$

where we have used

$$\int \sim |\psi|^{-\nu}$$

Thus $\omega = \nu(d-2)$

Scaling relations

$$\alpha = 2 - d\nu$$

$$\alpha + 2\beta + \gamma = 2$$

$$\gamma = \nu(2 - \eta)$$


From this, we obtain

$$\begin{aligned} 2 - d\nu + 2\beta + 2\nu - \eta\nu &= 2 \\ \Rightarrow \nu(d-2) &= 2\beta - \eta\nu \end{aligned}$$

$$\underline{\underline{\omega = 2\beta - \eta v}}$$

If $\eta > 0$, then $\omega < 2\beta$

d)



Comment: Searcher candidates used
 $2\beta - \omega = 2\beta - v(d-z)$

This is correct, but makes it less obvious if $2\beta > \omega$ or $2\beta < \omega$. One could have proceeded by:

$$\begin{aligned} \textcircled{x} \quad 2\beta - \omega &= 2\beta - v(d-z) \\ &= 2\beta - z + z - dv + 2v \\ &= \alpha + 2\beta - z + \delta + \eta v \\ &= \underbrace{\alpha + 2\beta + \delta - z}_{=0} + \eta v = \eta v \end{aligned}$$

⊗ Just stopping here, does not employ searchers

Problem 2

$$\langle \vec{J}(\vec{r}) \cdot \vec{J}(\vec{r}') \rangle = \left\langle \frac{e^{i\theta(\vec{r})} - e^{i\theta(\vec{r}')}}{2} \right\rangle$$

$$= e^{-\frac{1}{2K} \int \frac{d^d q}{(2\pi)^d} \frac{1 - \cos \vec{q} \cdot (\vec{r} - \vec{r}')}{q^2}}$$

$$\equiv I$$

$$H = \frac{c}{2} \int d^d x (\nabla \theta)^2$$

$$\text{Set } \vec{r}' = 0$$

$$\underline{d=3} \quad \vec{q} \cdot \vec{r} = qr \cos \theta$$

Do the angular integrations over φ and θ (azimuthal and polar angles), and $\cos(\vec{q} \cdot \vec{r}) = \cos \theta$

$$I = \frac{1}{K} \frac{1}{(2\pi)^3} 4\pi \int_{1/L}^{1/a} dq \left(\frac{1 - \sin q r}{q^2} \right)$$

a : Short-distance cutoff = 1

L : Long-distance cutoff

$$I = \frac{1}{K} \frac{2}{(2\pi)^2} \frac{1}{r} \int_{1/L}^{1/a} dx \left(\frac{1 - \sin x}{x} \right)$$

$$L \rightarrow \infty \Rightarrow$$

$$I = \frac{1}{2\pi^2 K} \frac{1}{r} \left\{ \frac{r}{a} - \text{Si}\left(\frac{r}{a}\right) \right\}$$

$$= \frac{1}{2\pi^2 K} \left[1 - \frac{1}{r} \text{Si}(r) \right]$$

$\text{Si}(x)$: Sine-integral

$$\text{Si}(x) = \int_0^x du \frac{\sin u}{u} \xrightarrow{x \rightarrow \infty} \frac{\pi}{2}$$

$$\lim_{r \rightarrow \infty} I(r) = \frac{1}{2\pi^2 K} - \frac{1}{2\pi^2 K} \left(1 - \frac{\text{Si}(r)}{r} \right)$$

$$\langle \vec{S}(\vec{r}) \cdot \vec{J}(0) \rangle = e$$

$$\xrightarrow{r \rightarrow \infty} e^{-\frac{1}{2\pi^2 K}} \neq 0$$

b)

$$\underline{\underline{\langle S(\vec{r}) \rangle = e^{-\frac{1}{4\pi^2 K}} \neq 0}}$$

Spin-waves do very little to
reduce $T=0$ magnetization in
3d FMS.

Problem 3

a) With the constraint removed the partition function reads

$$Z = \sum_{\{\vec{b}\}} e^{-\frac{1}{2K} \sum_{\vec{r}} \vec{b}^2}$$

$\vec{b} = (b_1, b_2, \dots, b_d)$ for d dimensions.

There are d b_i 's at every lattice site

$$Z = (Z_1)^{Nd} \quad N = \# \text{ lattice sites.}$$
$$Z_1 = \sum_{b=-\infty}^{\infty} e^{-\frac{1}{2K} b^2}$$

This Gaussian sum may be performed explicitly using the Jacobi θ -function

$$\psi_3(z, \tau) = \sum_{b=-\infty}^{\infty} g \frac{b^2}{b} \eta$$

$$g = e^{\pi i \tau} \quad ; \quad \eta = e^{2\pi i z}$$

$$g = e^{-\frac{1}{2K}} = e^{i\pi \tau} \Rightarrow i\pi \tau = -\frac{1}{2K}$$

$$\tau = \frac{i}{2\pi K}$$

$$Z_1 = \psi_3\left(0, \frac{i}{2\pi K}\right)$$

ψ_3 is an analytic function of its arguments, so Z_1 and hence $\ln Z_1$ and $\ln Z$ are also analytic functions of $T \Rightarrow$ No phase-transition.

$$b) \quad H = +J \sum_{\langle i,j \rangle} (1 - \cos(\theta_i - \theta_j)) - h \sum_i \cos \theta_i$$

Use a Villain-approximation
in both terms, to write

$$Z = \int D\theta \sum_{\{\vec{m}\}} \sum_{\{Q\}} e^{-\frac{K}{2} \sum_{\vec{p}, \mu} (\Delta_{\mu} \theta - 2\pi m_{\mu})^2 - \frac{\beta h}{2} \sum_{\vec{p}} (\theta - 2\pi Q)^2} \cdot e$$

$$K \equiv \beta J \quad ; \quad \omega \equiv \beta h$$

Next, use the identity

$$\int_{-\infty}^{\infty} dy e^{-\frac{a}{2} y^2 + ixy} \sim e^{-\frac{x^2}{2a}}$$

twice, to obtain

$$Z = \int D\theta \sum_{\{\vec{m}\}} \sum_{\{Q\}} \int D\vec{B} \int D\vec{R} e^{-\frac{1}{2K} \sum_{\vec{p}} \vec{B}^2 - \frac{1}{2\omega} \sum_{\vec{p}} R^2} e^{i \sum_{\vec{p}, \mu} B_{\mu} (\Delta_{\mu} \theta - 2\pi m_{\mu})} e^{i \sum_{\vec{p}} R (\theta - 2\pi Q)}$$

Collect terms involving e
(use $\sum_{\vec{r}} \vec{B} \cdot \Delta \vec{E} = - \sum_{\vec{r}} e (\vec{E} \cdot \vec{B})$)

to obtain a factor

$$e \sum_{\vec{r}} e (R - \vec{E} \cdot \vec{B})$$

Perform θ -integration $\Rightarrow$
at every lattice site, we have
a constraint

$$\vec{E} \cdot \vec{B} = R$$

Next, use Poisson summation
formula

$$\sum_{\vec{m}=-\infty}^{\infty} e^{-2\pi i \vec{m} \cdot \vec{B}} = \sum_{\vec{B}=-\infty}^{\infty} \delta(\vec{B} - \vec{B})$$

$$\sum_{Q=-\infty}^{\infty} e^{-2\pi i Q R} = \sum_{L=-\infty}^{\infty} \delta(L - R)$$


The partition function then becomes

$$Z = \sum_{\{\vec{B}\}} \sum_{\{\vec{L}\}} e^{-\frac{1}{2K} \sum_{\vec{r}} \vec{B}^2 - \frac{1}{2\omega} \sum_{\vec{r}} \vec{L}^2} \left(\prod_{\vec{r}} \delta_{\vec{\nabla} \cdot \vec{B}, \vec{L}} \right)$$

$\omega \neq 0$ allows for $\vec{L} \neq 0$

$$\vec{\nabla} \cdot \vec{B} = \vec{L} \neq 0$$

means that link-currents can start or end on a lattice point.



Such configurations have a non-zero probability of occurring

even at very high T .

One may "glue" together such small "dumbbells" to form large loops even at very high T .

In the superfluid language, one would say one has superfluidity even at very high T .

In the spin-language, one would say the system is ordered even at high T .

Finite h destroys the

order-disorder transition

In fact, h is allowed to take any value (integer)

Thus $\vec{\nabla} \cdot \vec{B} = 1$

is no constraint on $\vec{B}$!

This destroys the PT, since

$$Z = (Z_1, Z_2)^{Nd}$$

where Z_1 and Z_2

are Jacobi θ -functions.

c) $h \rightarrow 0 \Rightarrow \omega \rightarrow 0$

$$\frac{1}{2\omega} \rightarrow \infty$$

This suppresses all l 's
except $l=0$ at each
lattice site. Thus we have

$$Z = \sum_{\{\vec{B}\}} e^{-\frac{1}{2K} \sum_P \vec{B}^2} \left(\prod_P \delta_{\vec{\nabla} \cdot \vec{B}, 0} \right)$$

which is the loop-gas
representation of the XY-model.