

Department of Physics

Examination paper for FY3201 / 8902 Atmospheric Physics and Climate Change

Examination date: 1 June 2021

Examination time (from-to): 09:00-13:00

Permitted examination support material: All support materials are allowed

Academic contact during examination: Patrick Espy

Phone: +47 73 55 10 95

Technical support during examination: Orakel support services:

Link: innsida.ntnu.no/wiki/-/wiki/English/Orakel+Support+Services

Phone: +47 73 59 16 00

If you experience technical problems during the exam, contact Orakel support services as soon as possible before the examination time expires. If you don't get through immediately, hold the line until your call is answered.

OTHER INFORMATION

Only contact academic contact in case of errors or insufficiencies in the question set.

Saving: Answers written in Inspira are automatically saved every 15 seconds. If you are working in another program remember to save your answer regularly.

Cheating/Plagiarism: The exam is an individual, independent work. *Examination aids are permitted*. All submitted answers will be subject to plagiarism control. Read more about cheating and plagiarism here: innsida.ntnu.no/wiki/-/wiki/English/Cheating+on+exams

Notifications: If there is a need to send a message to the candidates during the exam (e.g. if there is an error in the question set), this will be done by sending a notification in Inspira. A dialogue box will appear. You can re-read the notification by clicking the bell icon in the top right-hand corner of the screen. All candidates will also receive an SMS to ensure that nobody misses out on important information. Please keep your phone available during the exam.

ABOUT SUBMISSION

Your answers will be submitted automatically when the examination time expires and the test closes, if you have answered at least one question. This will happen even if you do not click "Submit and return to dashboard" on the last page of the question set. You can reopen and edit your answer as long as the test is open. If no questions are answered by the time the examination time expires, your answers will not be submitted.

Withdrawing from the exam: If you wish to submit a blank test/withdraw from the exam, go to the menu in the top right-hand corner and click "Submit blank". This **can not** be undone, even if the test is still open.

Accessing your answer post-submission: You will find your answer in Archive when the examination time has expired.

Each incorrect or blank answer is 0 points. Answers have been randomized and are not exact. You must choose the closest answer!

You may take:

Molar mass of dry air: $\sim 29 \text{ kg/kmole}$

Molar mass of helium: $\sim 4 \text{ kg/kmole}$

Molar mass of H_2O : $\sim 18 \text{ kg/kmole}$

Molar mass of CO_2 : $\sim 44 \text{ kg/kmole}$

$N_A = 6.02 \times 10^{23} \text{ molecules/mole}$

Boltzmann's constant $k = 1.38 \times 10^{-23} \text{ J/K}$

$273 \text{ K} = 0^\circ\text{C}$ $1 \text{ hPa} = 10^2 \text{ Pa} = 10^2 \text{ N m}^{-2}$ $1 \text{ atm} = 1013 \text{ hPa}$ $g = 9.8 \text{ m s}^{-2}$ constant in z

Stefan–Boltzmann constant: $\sigma = 5.67 \times 10^{-8} \text{ W}\cdot\text{m}^{-2}\cdot\text{K}^{-4}$

Solar photospheric temperature, $T_s = 5786 \text{ K}$

Radius of the Sun = 695800 km

Radius of the Earth = 6370 km

1 AU (Earth-Sun distance) = $150 \times 10^6 \text{ km}$

Radius of Mercury = 6051 km

Mercury-Sun distance = 0.387 AU

Radius of Mars = 3396 km

Mars-Sun distance = 1.52 AU

Latent heat of vaporization water: $L_v = 2.5 \times 10^6 \text{ J}\cdot\text{kg}^{-1}$

Density of liquid water = $1000 \text{ kg}\cdot\text{m}^{-3}$

Latent heat of sublimation ice: $L_i = 2.8 \times 10^6 \text{ J}\cdot\text{kg}^{-1}$

Density of water vapour = $5 \times 10^{-3} \text{ kg}\cdot\text{m}^{-3}$

Gas constant for water vapour: $R_v = 461 \text{ J}\cdot\text{K}^{-1}\cdot\text{kg}^{-1}$

Surface tension of water droplet $75 \times 10^{-3} \text{ N}\cdot\text{m}^{-1}$

Values for dry air:

$C_p = 1004 \text{ J}\cdot\text{K}^{-1}\cdot\text{kg}^{-1}$

$C_v = 718 \text{ J}\cdot\text{K}^{-1}\cdot\text{kg}^{-1}$ $R_d = 287 \text{ J}\cdot\text{K}^{-1}\cdot\text{kg}^{-1}$

$\gamma = C_p / C_v$

$\kappa = R_d / C_p$

$R_d = C_p - C_v$

$\Gamma_{da} = 9.8 \text{ K/km}$

Clausius–Clapeyron relation: $e_s = 6.112 \text{ hPa} \cdot \exp\left[\frac{L_v}{R_v}\left(\frac{1}{273 \text{ K}} - \frac{1}{T}\right)\right]$

Some integrals that may be of use:

$$\int x^m e^{(ax)} dx = \frac{x^m e^{(ax)}}{a} - \frac{m}{a} \int x^{(m-1)} e^{(ax)} dx$$

$$\int x e^{(ax)} dx = \frac{e^{(ax)} (ax - 1)}{a^2}$$

$$\text{For } a > 0 \quad \int_0^\infty e^{(-ax)} dx = \frac{1}{a}$$

$$\int \frac{1}{a + bx} dx = \frac{\ln(a + bx)}{b}$$

1. Geostrophic flow occurs when the wind blows

- (1) parallel to the isobars or contours or pressure.
- (2) in the direction of the Coriolis force.
- (3) in the direction of the pressure gradient force.
- (4) at an angle between 10 and 30 to the contours and towards low pressure.
- (5) at an angle between 10 and 30 to the contours and towards high pressure.

In geostrophic flow, the pressure gradient force is balanced by the Coriolis force. When this happens, the wind ceases to change, and results in the wind blowing parallel to the isobars.

2. Of the gases listed below, which is NOT believed to be responsible for enhancing the earth's greenhouse effect?

- a. molecular oxygen (O₂)
- b. chlorofluorocarbons (CFCs)
- c. nitrous oxide (N₂O)
- d. carbon dioxide (CO₂)
- e. methane (CH₄)

A greenhouse gas has high transmission in the visible, but absorbs appreciably in the infrared. Since infrared wavelengths interact with molecular vibrations and rotations, the molecule must possess a permanent dipole moment, or one that is induced by vibration or rotation. O₂ is a symmetric (homonuclear) molecule that does not possess a permanent dipole moment. Since the only vibrations are along the internuclear axis, one cannot be induced. Hence it cannot interact with infrared light.

3. The most abundant gas in the stratosphere is:

- a. nitrogen (N₂).
- b. oxygen (O₂).
- c. carbon dioxide (CO₂).
- d. ozone (O₃).
- e. chlorofluorocarbons (CFCs).

Ozone is never the major species, but always a minor one. In the well-mixed atmosphere is N₂ always the major species.

4. How do you find the Lifting Condensation Level (LCL) on a Skew-T diagram?

- a. Find the intersection between a parcel's dry adiabat and constant $\mu_s = \mu$ of the parcel
- b. Find a parcel's water vapour mixing ratio using its dew point temperature.
- c. Find the region where the temperature starts to rise with altitude.
- d. Find where the atmospheric lapse rate is equal to the dry adiabatic lapse rate.

The LCL is where the water vapour contained in the parcel, μ , may begin to condense. The parcel will therefore follow the dry adiabat until it reaches the point where $\mu_s = \mu$. This point can then be found by tracing the isopleth of $\mu_s (= \mu)$ until it intersects the parcel's dry adiabat.

5. If the earth's average surface temperature were to increase, the amount of radiation emitted from the Earth's surface would ____, and the wavelength of peak emission would shift toward ____ wavelengths.

- a. increase; shorter
- b. increase; longer
- c. decrease; shorter
- d. decrease; longer

The total energy emitted is found from Stefan-Boltzmann relation $F(w/m^2) = \sigma_{SB} * T^4$, where σ_{SB} is a constant. The peak wavelength of an emission is given by Wein displacement law $\lambda_{peak} = \text{Constant}/T$. Therefore, an increase in temperature will result in a shorter wavelength of peak emission.

6. a) A spherical drop of water of radius 20 nm is stable (neither growing nor shrinking) at a temperature of 15 C. what is the relative humidity around the droplet?

For a drop, one must use Kelvin's formula where ρ_L = density of liquid water, 1000 kg/m³, R_v is the specific gas constant for water vapour, 461 J/K/kg, and r is the radius of a stable drop (the turning point is the free energy equation)

$$e = e_s \cdot e^{\left(2 \frac{\gamma}{\rho_L R_v T r}\right)}$$

Here, γ is the surface tension = 75x10⁻³ kg/m. The equilibrium or saturated partial pressure of water vapour over a flat surface of water, e_s , is given as

$$e_s = 6.112 \cdot e^{\left(\frac{L_v \left(\frac{1}{273} - \frac{1}{T}\right)}{R_v}\right)}$$

Where the latent heat of vaporization, $L_v = 2.5 \times 10^6$ J/kg. Substituting in these values for $T = (15 + 273)K$ and a droplet radius of 20×10^{-9} m gives: $e_s = 17.63$ hPa, and $e = 18.66$ hPa.

The relative humidity is therefore $e/e_s = 106\%$

b) An airplane economy cabin is pressurized to 818 hPa with a temperature 20 C and a relative humidity of 15% when flying at 11 km. What is the dew point temperature in the cabin?

The dew point temperature is the temperature where the partial pressure of water vapour in the air, e , is equal to the saturation vapour pressure, $e_s(T_d)$. The saturation vapour pressure is as above, and if we substitute $e = e_s$, and $T = T_d$, we can solve for T_d

$$T_d = 273. \frac{L_v}{L_v - 273. \ln(.1636125654 \cdot e) R_v}$$

Where L_v and R_v are given as above. We still need e , but this is given by $RH=e/e_s(T)$, where $e_s(T)$ is, as before:

$$e_s = 6.112 \cdot e^{\left(\frac{L_v \left(\frac{1}{273} - \frac{1}{T} \right)}{R_v} \right)}$$

At a temperature of $T=(20+273)K$, the saturated partial pressure of water vapour is $e_s=24.50$ hPa. With a relative humidity of 15%, the partial pressure of water vapour, $e = 0.15 \cdot 24.5$ hPa = 3.67 hPa. Given this, we can calculate the dew point temperature to be: $T_d = 266.3$ K = -6.66 C.

c) A person perspires. How much liquid water (as a percentage of the person's mass) must evaporate to lower the temperature of the person by 5.0 C. Take the specific heat of the human body to be that of water, $C_{pw} = 4200$ J/kg/K.

Evaporating a mass of water, M_w , will release an amount of heat, $dQ = L_v \cdot M_w$ into the environment

Losing an amount of heat, dQ , from a body of mass M_p with a heat capacity $C_{pw} = 4200$ J/kg/K will drop its temperature by $dQ = C_{pw} \cdot M_p \cdot dT$

Equating the dQ 's and solving for $M_w/M_p = dT \cdot C_{pw} / L_v$

For $dT=5$ K (remember a change in temperature is the same in K as in C) and the values for C_{pw} and L_v we get that $M_w/M_p = 0.0075 = 0.75\%$

7. An exoplanet orbits its star at a distance $R_{orbit} = 0.41$ AU, and has a radius $R_p = 1.34 \cdot R_{earth}$. The planet has an albedo of 0.4 and emissivity of 1.

The star it orbits has a radius $R_{st} = 0.6 \cdot R_{sun}$ and a photosphere blackbody temperature of $T_{st} = 4400$ K

a. What is the planet's equilibrium temperature assuming it has no atmosphere?

We need to know the "stellar" constant for this planet. that is the W/m^2 it puts on the planet. The irradiance W/m^2 at the photosphere of the star is given (with $\sigma_{SB} =$ Stefan Boltzmann constant):

$$F_{st} = \sigma_{SB} \cdot T_{st}^4$$

The star's total power in watts is this times the area of the photosphere. This is the luminosity of the point source radiating in all directions ($4\pi sr$).

$$L_{st} = 4\pi \cdot R_{st}^2 \cdot \sigma_{SB} \cdot T_{st}^4$$

At the planet, this luminosity is distributed over a sphere $4\pi \cdot (R_{orbit})^2$ giving the planet's stellar constant as:

$$F_p = \frac{R_{st}^2}{R_{orbit}^2} \cdot \sigma_{SB} \cdot T_{st}^4$$

We can calculate this intermediate step and find that the stellar constant for this planet is 979.3 W/m^2

Now the planet absorbs (1-albedo) of this stellar flux over its cross-sectional area of $\pi \cdot (R_p)^2$, and radiates over its surface area $= 4\pi \cdot R_p^2$ as $\epsilon \cdot \sigma_{SB} \cdot (T_p)^4$

At equilibrium, energy in = energy out and the radius of the planet cancels:

$$\pi \cdot R_p^2 \cdot F_p (1 - a) = 4\pi \cdot R_p^2 \cdot \epsilon \cdot \sigma_{SB} \cdot T_{st}^4$$

We can solve for T_{st} and find:

$$T_p = \frac{1}{2} \frac{\sqrt{2} (-F_p (-1 + a) \epsilon^3 \sigma_{SB}^3)^{\left(\frac{1}{4}\right)}}{\epsilon \sigma_{SB}}$$

Using the values given, $T_p = 225.6 \text{ K} = -47.4 \text{ C}$

b. At what wavelength does the radiance of this star with a blackbody temperature of 4400 K peak?

For this we can use Wiens displacement law for a stellar temperature of 4400 K, the peak wavelength in microns is:

$$\lambda = 2898 \frac{1}{T_{st}}$$

$$\lambda = 0.6586 \mu = 658.6 \text{ nm}$$

c. Another exoplanet in the same system has a stellar flux constant F_p of 800 W/m^2 at its orbit, The planet has an albedo of 0.4, and emissivity of 1, but there is a compound in the atmosphere that passes all infrared light but absorbs in the visible.

A probe has told us that the temperature of the surface of this planet is -85 C . What is the short wavelength transmission of the atmosphere of this planet?

We could re-derive the radiative equilibrium equation with an atmosphere, but if we take the short and long wavelength transmissions to be TS and TL , we know from the class notes that a planet at temperature T_p , with albedo a and stellar flux F_p , has a radiative balance equation given by:

$$\epsilon \sigma T_p^4 = \frac{1}{4} \frac{(1 - a) F_p (1 + TS)}{1 + TL}$$

We can solve for TS ,

$$TS = \frac{4(1 + TL) \cdot \epsilon \cdot \sigma_{SB} \cdot T_p^4}{(1 - a) \cdot F_p} - 1$$

And find that $TS = 0.18 = 18\%$

8. Sunlight coming into the Earth's atmosphere can be absorbed and heat the atmosphere.

The Sun is at a solar ZENITH angle (angle of the Sun from overhead) of $\chi = 45$ degrees. You may assume no scattering, and take the optical depth of incoming solar radiation (averaged over all wavelengths), the density of air, and the mass mixing ratio of the absorber to be defined as:

$$\tau(Z) = \int_Z^\infty \rho(z) \mu(z) k dz$$

$$\rho(z) = \rho_0 e^{\left(-\frac{z}{H_m}\right)}$$

$$\mu(z) = \mu_0 e^{\left(-\frac{z}{H_v}\right)}$$

Where $\rho(z)$ is the mass density of air, $\mu(z)$ is the mass mixing ratio of the component in the air that attenuates the radiation, and k is the mass absorption coefficient.

The air density at the surface, $\rho_0 = 1.2 \text{ kg/m}^3$ and the scale height of air is $H_m = 7 \text{ km}$. In addition, $\mu_0 = 0.01 \text{ (kg of absorbing component)/(kg air)}$, and its scale height $H_v = 4 \text{ km}$. Finally, the mass absorption coefficient, k is constant and $k = 0.11 \text{ m}^2/(\text{kg of absorbing compound})$.

a) Calculate the optical depth τ and its derivative with height $d\tau/dz$ at a height of 5 km in the atmosphere

So, the first thing is to substitute in for ρ and μ into the integral

$$\tau = \int_0^{\infty} \rho_0 e^{\left(-\frac{z}{H_m}\right)} \mu_0 e^{\left(-\frac{z}{H_v}\right)} k dz$$

This is not too hard to integrate as the exponents of the exponentials add, and can be factored out. With k a constant, we get:

$$\tau = \frac{\rho_0 \mu_0 k H_m H_v e^{\left(-\frac{z(H_v + H_m)}{H_m H_v}\right)}}{H_v + H_m}$$

and of course, its derivative

$$\frac{d\tau}{dz} = -\rho_0 \mu_0 k e^{\left(-\frac{z(H_v + H_m)}{H_m H_v}\right)}$$

Substituting in the values given yields:

$$\tau = 0.47, \text{ and } \frac{d\tau}{dz} = -0.00019 \text{ m}^{-1} = -0.19 \text{ km}^{-1}$$

b) If the incoming solar flux at the top of the atmosphere is 400 W/m^2 , and the solar ZENITH angle, χ , is 45 degrees, calculate the NET UPWARD flux of solar radiation at 7 km if the optical depth is 0.2 at this altitude

First things first. The ZENITH angle vs $\mu = \cos(\theta)$ is shown here:

As you can see $\cos(\chi) = -\mu$. Therefore, we can use the equation for downward flux from the short wavelength heating lecture, but the one for χ instead of μ . This is given by:


$$F_v^{\downarrow}(z) = -F_{v\infty}^{\downarrow} \cdot \mu \cdot e^{\frac{\tau(z)}{\mu_s}} = F_{v\infty}^{\downarrow} \cdot \cos(\chi_s) \cdot e^{-\frac{\tau(z)}{\cos(\chi_s)}}$$

Where $F_{v\infty}^{\downarrow} = 400 \text{ W} \cdot \text{m}^{-2}$

All we have to do now is pay attention to the fact that we want the net UPWARD flux, which is:

$$F_v(z) = [F_v^{\uparrow}(z) - F_v^{\downarrow}(z)] = F_{v\infty}^{\downarrow} \cdot \mu \cdot e^{\frac{\tau(z)}{\mu_s}} = -F_{v\infty}^{\downarrow} \cdot \cos(\chi_s) \cdot e^{-\frac{\tau(z)}{\cos(\chi_s)}}$$

Putting in the numbers, we get the net upward flux $F_v(z=7\text{km}) = -213 \text{ W/m}^2$.



c) Calculate the heating rate in Kelvin/day due to short wavelength radiation at 3 km if the downward solar flux at the top of the atmosphere is 400 W/m^2 .

The solar ZENITH angle is $\chi = 45$ degrees, and at 3 km, the downward solar flux is 70 W/m^2 , the density is $\rho = 0.78 \text{ kg/m}^3$, the optical depth is $\tau = 1$ and its derivative, $d\tau/dz$, is -0.4 km^{-1} .

Well, the heating rate in W/kg, when divided by the specific heat C_p in J/kg/K, will give the heating rate in K/sec. That heating rate is given in terms of the net upward flux as:

$$\frac{\partial}{\partial t} Q = - \frac{\frac{\partial}{\partial z} F_v(z)}{\rho(z) C_p}$$

And in the previous part (b.) we have an expression for $F_v(z)$:

$$F_v(z) = -F_{v\infty}^{\downarrow} \cdot \cos(\chi_s) \cdot e^{-\frac{\tau(z)}{\cos(\chi_s)}}$$

When we differentiate this with respect to z , we get:

$$\frac{\partial}{\partial t} Q = - \frac{-F_{v\infty}^{\downarrow} \left(\frac{\partial}{\partial z} \tau(z) \right) e^{-\frac{\tau(z)}{\cos(\chi_s)}}}{\rho(z) C_p}$$

Now, kindly old professor Espy actually gave you everything you need to calculate this without differentiating τ . That's because you needed to do that in part a. and if you didn't, then you shouldn't lose points twice! Plugging in the numbers, you get:

$$\frac{dQ}{dt} = 0.00005 \frac{K}{s} = 0.003 \frac{K}{min} = 0.17 \frac{K}{hr} = 4 \frac{K}{day}$$

d) In an isothermal atmosphere of temperature 220 K, at what pressure level will the Doppler-line width be the same as the collision-line width at a wavelength of 15 microns.

We can use the formula for line shape as a function of pressure and temperature given in lecture

$$\gamma_L = 0.01 \text{ cm}^{-1} \cdot \left(\frac{p}{1000 \text{ hPa}} \right) \left(\frac{273 \text{ K}}{T} \right)^{1/2}$$

$$\gamma_D = 0.003 \text{ cm}^{-1} \cdot \left(\frac{\nu}{1000 \text{ cm}^{-1}} \right) \left(\frac{T}{300 \text{ K}} \right)^{1/2}$$

Equate them and solve for P . We of course need to turn 15 μ into wavenumbers in cm^{-1} . That comes out to be $\nu = 1/(15 \times 10^{-6} \text{ m} \cdot 100 \text{ cm/m}) = 666.7 \text{ cm}^{-1}$.

If we equate γ_L and γ_D , and solve for P , we get:

$$P = 0.3 \frac{\nu \cdot T}{\sqrt{273 \cdot 300}}$$

At 666.7 cm^{-1} and 220 K, we get a pressure $P=154 \text{ hPa}$

9. On Venus $g=8.87 \text{ m/s}^2$, the specific gas constant for dry air is $R_{dv} = 195.5 \text{ J/K/kg}$, and the specific heat at constant pressure, $C_p = 846 \text{ J/K/kg}$.

The temperature profile is given by $T=T_0 \cdot \exp(-a \cdot Z)$, with $T_0=783\text{K}$ and $a=1.68\text{e-}5 \text{ m}^{-1}$

The surface pressure is $P_0 = 93320.33 \text{ hPa}$, and the atmosphere behaves like a dry ideal gas.

a. What is the buoyancy period at 59 km

The Brunt frequency, or buoyancy frequency, in radians/sec is given by:

$$N = \sqrt{\frac{g \left(\frac{g}{C_p} + \frac{dT}{dz} \right)}{T}}$$

Of course, we have a different g , C_p , and we need to calculate $T(z=59 \text{ km})$ and dT/dz there. We are asked for a period, and the brunt period is given by:

$$T_{\text{Brunt}} = \frac{2\pi}{N} \text{ in seconds}$$

Putting in the values for Venus, and calculating the temperature and temperature gradient:

$$T = T_0 e^{(-a \cdot z)} \text{ and } \frac{dT}{dz} = -T_0 \cdot a \cdot e^{(-a \cdot z)}$$

At 59 km, we get values of $T=291 \text{ K}$, $\Gamma_d=g/C_p=0.01\text{K/m}$, $\Gamma_{\text{env}}=-\frac{dT}{dz}=0.005 \text{ K/m}$, giving

$N=0.013 \text{ rad/sec}$ and $T_{\text{Brunt}}=480.5 \text{ s} = 8 \text{ minutes}$

Just as a note, at about 59 km, Venus has a temperature and pressure about the same as at the surface of the Earth!

b. Is the atmosphere stable with respect to vertical motions at 59 km?

It is stable, because $N^2 > 0$, so that a displaced parcel will return to its original position, oscillating under the influence of gravity and buoyancy. Some answered conditionally stable or conditionally unstable. No information is given from which one could deduce this. However, if the 2 points from this answer changed the grade then it was given.

c. Is the atmosphere stable with respect to vertical motions at the surface?

Here we have to calculate N^2 for the surface conditions. Now we find that at the surface, we get values of $T=783 \text{ K}$, $\Gamma_d=g/C_p=0.01\text{K/m}$, $\Gamma_{\text{env}}=-\frac{dT}{dz}=0.013 \text{ K/m}$, giving $N^2=-0.00003 \text{ rad/s}$. So *it is unstable* since N would be imaginary, and a displaced parcel would continue in the direction of the displacement

d. What is the Scale Height, H , at 59 km and at the surface?

Hopefully, you did not have the same brain fart that I did during the spørretime and remembered that:

$$H = \frac{RT}{g} = \frac{RT_0 e^{(-a \cdot z)}}{g}$$

Where R is now R_{av} and g is appropriate for Venus, at *59 km, this gives $H=6.4 \text{ km}$* , and at the *surface $H=17.3 \text{ km}$* .

e. What is the pressure at 59 km?

The hydrostatic equation is $\frac{dP}{dz} = -g \cdot \rho$. Substituting in for ρ using the ideal gas law, $P=\rho RT$, we get that:

$$\frac{dP}{P} = -\frac{g}{R \cdot T(z)} \cdot dz$$

Or

$$P = P_0 e^{\left(\int_0^Z \frac{g}{R T(z)} dz \right)}$$

But we know that $T(z)$ is, $T(z) = T_0 e^{(-az)}$, so we need to integrate:

$$\int_0^Z -\frac{g e^{(az)}}{R T_0} dz = -\frac{g (e^{(aZ)} - 1)}{R T_0 a}$$

So, now P is:

$$P = P_0 e^{\left(-\frac{g (e^{(aZ)} - 1)}{R T_0 a} \right)}$$

Putting in the appropriate values (including $R=R_{dv}$), we get $P=27072 \text{ Pa} = 270 \text{ hPa} = 27 \text{ kPa}$

f. If a parcel of air at the surface, where the pressure is $P_0=9332033 \text{ Pa}$, is moved adiabatically to an altitude of 700 m where the pressure is $0.96 \cdot P_0$, what is the parcel's final temperature?

This is our old friend the potential temperature, and the temperature T at pressure P is:

$$Tp = \frac{\theta}{\left(\frac{P_0}{P} \right)^\kappa}$$

Where θ is the temperature at our reference pressure, P_0 . We do need to calculate the new value for κ on Venus, however. That would be $\kappa=R_{dv}/C_{pV} = 0.23$.

If we choose our reference pressure at the surface, where we know the temperature is $T_0 = \theta$, we can calculate the temperature of the parcel at $P = 0.96 P_0$. We get that $Tp=776 \text{ K}$.

Note, the atmospheric temperature at 700 m is 773 K, so cooler than the parcel. Thus the atmosphere is unstable with regard to vertical motion.

10. Sulfur dioxide, (SO_2) is a pollutant released in combustion of coal and other high-sulfur fuels that can cause acid rain. Although Norway produces very little, it is transported by the prevailing winds to Norway from large industrial regions in Europe as well as from volcanic sources in Iceland. Thus, its concentration in Norway is highly variable between 1 and 25 ppbv (parts per billion by volume).

a. At 1000 hPa and 22 C, the mixing ratio of SO_2 measured one day in Bergen was 5 ppbv. What was the mass per unit volume of SO_2 ?

We will need the number of SO_2 molecules and multiply by the weight of each molecule. To get the number of molecules, we can go back to the definition of the mixing ratio: $v = \frac{N_{\text{SO}_2}}{N_{\text{total}}}$, and we can obtain

N_{total} from the ideal gas law: $N_{\text{total}} = \frac{P}{kT}$, where k is Boltzmann's constant. Thus, $N_{\text{SO}_2} = \frac{vP}{kT}$. We have everything we need with $P=100000 \text{ Pa}$, $T=22+273=295\text{K}$, and $v=5\text{e-}9$ giving:

$$N_{\text{SO}_2} = 1.23 \times 10^{17} \text{ molecules} \cdot \text{m}^{-3}.$$

We can verify this yields the correct mixing ratio by calculating $N_{\text{total}}=2.46 \times 10^{25}$ giving

$$v = \frac{1.23 \times 10^{17}}{2.46 \times 10^{24}} = 5 \text{ ppb}_v$$

Next, we need the mass of an SO_2 molecule. From the periodic table, this is $32+2 \cdot 16=64 \text{ g} \cdot \text{mole}^{-1}$, and there are $6.02 \times 10^{23} \text{ molecules} \cdot \text{mole}^{-1}$. This means the mass per unit volume, or density, ρ , is:

$$\rho_{\text{SO}_2} = \frac{1.23 \times 10^{17} \text{ molecules} \cdot \text{m}^{-3}}{6.02 \times 10^{23} \text{ molecules} \cdot \text{mole}^{-1}} \cdot 64 \text{ g} \cdot \text{mole}^{-1} \cdot 10^{-3} \text{ kg} \cdot \text{mole}^{-1}$$

This gives a density of $\rho_{\text{SO}_2} = 1.31 \times 10^{-8} \text{ kg} \cdot \text{m}^{-3} = 13.1 \mu\text{g} \cdot \text{m}^{-3}$

b. If a mass of $50 \mu\text{g}/\text{m}^3$ of SO_2 is entirely converted into sulfuric acid, H_2SO_4 , what is the resulting mass per volume of sulfuric acid at 1000 hPa and 22 C?

We could start over, or recognize that:

The number of sulphur atoms S = number of SO_2 molecules = number of H_2SO_4 molecules.

Instead of each S atom being associated with a $64 \text{ g}/\text{mole}$ SO_2 molecule, it is associated with a molecule of $\text{H}_2\text{SO}_4 = 1 + 32 + 4 \cdot 16 = 98 \text{ g}/\text{mole}$. We can take the $50 \text{ micrograms}/\text{m}^3$ of SO_2 , multiply by the ratio of molecular weights: $98/64$ to get mass H_2SO_4 :

$$\rho_{\text{H}_2\text{SO}_4} = 50 \times 10^{-9} \text{ kg} \cdot \text{m}^{-3} \cdot \frac{98}{64} = 7.66 \times 10^{-8} \text{ kg} \cdot \text{m}^{-3} = 76.6 \mu\text{g} \cdot \text{m}^{-3}$$

c. Originally, the Martian atmosphere was thought to consist only of 95% CO_2 , 3% N_2 and 2% Ar. However, recently, it was discovered that it also contains 0.2% O_2 . By how much did the initial estimate of the mean molecular weight of the Martin atmosphere change when this additional species, O_2 was taken into account.

The molecular weight of a mixture of gases is the sum of the products of the mixing ratios of each component times its molar mass $M_m = \sum v_i \cdot M_{mi}$. For Mars, we can get the molar masses from the periodic table where we have that M_m of $\text{CO}_2=44$, $\text{N}_2 = 28$, Ar, = 40, and for later, $\text{O}_2 = 32$ all in g/mole .

The molecular weight of the Martian atmosphere without O_2 is therefore:

$$M_{m_{\text{no O}_2}} = 0.95 \cdot 44 + 0.03 \cdot 28 + 0.02 \cdot 40 = 43.44 \text{ g} \cdot \text{mole}^{-1}$$

Now, if we replace 0.2% of this mixture with O_2 , we calculate the new molecular weight by:

$$M_{m_{\text{O}_2}} = (1 - v_{\text{O}_2}) \cdot M_{m_{\text{no O}_2}} + v_{\text{O}_2} \cdot M_{\text{O}_2} = (1 - 0.002) \cdot 43.44 + 0.002 \cdot 32 = 43.41 \text{ g} \cdot \text{mole}^{-1}$$

Thus, the molecular mass has gone down, **decreasing by $0.03 \text{ kg}/\text{kmole}$**