

NTNU Trondheim, Institutt for fysikk**Examination for FY3452 Gravitation and Cosmology**

Contact: M. Kachelrieß, tel. 99890701

*Allowed tools: all***1. Gravitational waves.**

a.) How many independent components has a metric perturbation $h_{\mu\nu}$ described by the wave equation in the harmonic gauge: (2 pt)

- ☐ 2
☐ 4
☐ 5
☐ 6
☐ 10
☐ 16

b.) Consider the gravitational wave produced by a binary system of two equal masses M on a circular orbit in the xy plane which is seen by three observers at large distance on the x axis, the y axis and the z axis. Determine the observed polarisation by expressing the wave as $h_{\mu\nu} = \sum_a h^{(a)} \varepsilon_{\mu\nu}^{(a)}$, where $\varepsilon_{\mu\nu}^{(a)}$ is an appropriate basis of the polarisation states. (12 pts)

2. Schwarzschild black hole.

The Riemann tensor in Schwarzschild coordinates is

$$R_{0101} = \frac{2M}{r^3}, \quad R_{0202} = R_{0303} = -\frac{M}{r^3}, \quad (1)$$

$$R_{1212} = R_{1313} = \frac{M}{r^3}, \quad (2)$$

$$R_{2323} = -\frac{2M}{r^3}, \quad (3)$$

all other elements which cannot be obtained by its (anti-) symmetry properties are zero.

a.) Show that the Riemann tensor in the inertial system of a freely falling observer has the same form as given above. (8 pts)

b.) The distance n^i of two freely falling particles changes as

$$\ddot{n}^i = R^i_{00j} n^j.$$

Consider now a cube with mass m assumed to be a rigid body of length L . What are the forces and the stresses acting on the cube at the distance r ? [Hint: Consider in (a Newtonian picture) the force which has to counter-balance the gravitational acceleration

of a mass element dm of the rigid body.] (16 pts)

c.) Evaluate numerically the stress for values appropriate for a human. (6 pts)

3. Cosmology.

The static Einstein universe contains no radiation and has positive curvature.

a.) For a given value of Λ , there is a unique value of the matter density ρ_0 such that $\ddot{R} = 0$. Express ρ_0 and the scale factor R_0 through Λ . (6 pts)

b.) Consider a small perturbation, $\rho_m = \rho_0 + \delta\rho$. Use the Friedmann equation to show that the resulting change δR satisfies

$$\frac{d^2\delta R}{dt^2} = B\delta R \quad (4)$$

with B as constant. Is the Einstein universe stable or not? (10 pts)

4. Symmetries.

a.) Consider in Minkowski space a complex scalar field ϕ with Lagrange density

$$\mathcal{L} = s_1 \frac{1}{2} \partial_\mu \phi^\dagger \partial^\mu \phi + s_2 \frac{1}{4} \lambda (\phi^\dagger \phi)^2.$$

Name the symmetries of the Lagrangian and explain your choice for the signs s_1 and s_2 . (6 pts)

b.) Consider in Minkowski space the following Lagrange density

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - j^\mu A_\mu,$$

where A_μ is the photon field, $F_{\mu\nu}$ the field-strength tensor, and j^μ an external current. Calculate the resulting change of the action under a gauge transformation $A_\mu \rightarrow A_\mu + \partial_\mu \Lambda$. Show that the action is invariant, if the current is conserved. (6 pts)

c.) Generalise now the two Lagrangians to a general space-time. Explain the general rules you apply. Is the procedure unique? (4 pts)