



Solution to the exam in
FY3464 QUANTUM FIELD THEORY I
 Wednesday october 17, 2007

This solution consists of 4 pages.

Problem 1.

Consider the model defined by the Lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - \frac{1}{2} m^2 \varphi^2 + \lambda \varphi \mathbf{E} \cdot \mathbf{B}, \quad (1)$$

where φ is a real scalar field, $\mathbf{E} = -\dot{\mathbf{A}}$, and $\mathbf{B} = \nabla \times \mathbf{A}$.

- a) Find the canonically conjugate field Π_φ of φ .

$$\Pi_\varphi = \frac{\partial \mathcal{L}}{\partial \dot{\varphi}} = \dot{\varphi}. \quad (2)$$

- b) Find the canonically conjugate field $\Pi_{\mathbf{A}}$ of \mathbf{A} .

$$\Pi_{\mathbf{A}} = \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{A}}} = -\frac{\partial \mathcal{L}}{\partial \mathbf{E}} = -\lambda \varphi \mathbf{B}. \quad (3)$$

- c) Find the Hamiltonian density \mathcal{H} of this model.

$$\mathcal{H} = \Pi_\varphi \dot{\varphi} + \Pi_{\mathbf{A}} \cdot \dot{\mathbf{A}} - \mathcal{L} = \frac{1}{2} \Pi_\varphi^2 + \frac{1}{2} \nabla \varphi \cdot \nabla \varphi + \frac{1}{2} m^2 \varphi^2. \quad (4)$$

- d) We use natural units. What is the mass dimension of the coupling parameter λ :

- (i) In 4 space-time dimensions? (ii) In d space-time dimensions?

By comparing dimensions between the first two terms in the Lagrangian (1),

$$[m^2 \varphi^2] = [m]^2 [\varphi]^2 = [\partial_\mu \varphi \partial^\mu \varphi] = \ell^{-2} [\varphi]^2,$$

we note that mass and length dimensions are inverse in natural units (as is also obvious from the expression for Compton wavelength). From the fact that the action $S = \int d^d x \mathcal{L}$ must be dimensionless (dimension of \hbar) it follows that \mathcal{L} must have dimension $\ell^{-d} = [\partial_\mu \varphi \partial^\mu \varphi] = [\varphi]^2 \ell^{-2}$, i.e. that

$$[\varphi] = \ell^{(2-d)/2} = [m]^{(d-2)/2}.$$

The dimension of $\mathbf{E} \cdot \mathbf{B}$ cannot be read out of the Lagrangian (1), although it can be seen that $[\mathbf{E}] = [\mathbf{B}]$ from their relations to \mathbf{A} . This was an oversight in the exam set; it was assumed that the dimensions are the same as in the “standard” case, when there is a term $\mathcal{L}_{\text{Maxwell}} = \frac{1}{2}(\mathbf{E}^2 - \mathbf{B}^2)$ in the Lagrangian. This leads to the result that that $[\mathbf{E}] = [\mathbf{B}] = m^{d/2}$. It now follows that

$$[\lambda] = [m]^d [\varphi]^{-1} [\mathbf{E} \cdot \mathbf{B}]^{-1} = [\varphi]^{-1} = [m]^{(2-d)/2}. \quad (5)$$

I.e, the mass dimension is -1 in the case when $d = 4$.

e) Find the Euler Lagrange equation for φ .

We find that

$$\frac{\partial \mathcal{L}}{\partial \partial_\mu \varphi} = \partial^\mu \varphi \quad \text{and} \quad \frac{\partial \mathcal{L}}{\partial \varphi} = -m^2 \varphi + \lambda \mathbf{E} \cdot \mathbf{B},$$

so that the Euler Lagrange equation,

$$\partial_\mu \frac{\partial \mathcal{L}}{\partial \partial_\mu \varphi} = \frac{\partial \mathcal{L}}{\partial \varphi},$$

becomes

$$(\square + m^2) \varphi = \lambda \mathbf{E} \cdot \mathbf{B}. \quad (6)$$

f) Find the Euler Lagrange equation for \mathbf{A} .

It is convenient to first write

$$\mathbf{E} \cdot \mathbf{B} = -\varepsilon^{\ell j k} \dot{A}^\ell \partial_j A^k = -\varepsilon^{j k \ell} \dot{A}^j \partial_k A^\ell,$$

so that we find

$$\frac{\partial \mathcal{L}}{\partial \dot{A}^\ell} = -\lambda \varphi \varepsilon^{\ell j k} \partial_j A^k = -\lambda \varphi B^\ell \quad \text{and} \quad \frac{\partial \mathcal{L}}{\partial_k A^\ell} = -\lambda \varphi \varepsilon^{j k \ell} \dot{A}^j.$$

Thus, the Euler-Lagrange equation becomes

$$-\lambda \left[\partial_0 (\varphi B^\ell) - \varepsilon^{\ell k j} \partial_k (\varphi \dot{A}^j) \right] = -\lambda \left(\dot{\varphi} B^\ell + \varepsilon^{\ell k j} E^j \partial_k \varphi \right) - \lambda \varphi \left[\dot{B}^\ell - (\nabla \times \dot{\mathbf{A}})^\ell \right].$$

The last term on the right vanishes. Assuming $\lambda \neq 0$ we arrive at the equation

$$\mathbf{B} \dot{\varphi} - \mathbf{E} \times \nabla \varphi = 0. \quad (7)$$

Comment: By introducing the electromagnetic field tensor $F_{\mu\nu} = \partial_\nu A_\mu - \partial_\mu A_\nu$, and its dual field tensor $\tilde{F}^{\mu\nu} = \frac{1}{2} \varepsilon^{\mu\nu\lambda\sigma} F_{\lambda\sigma}$, this equation can be written in manifestly covariant form¹,

$$\tilde{F}^{\mu\nu} \partial_\nu \varphi = 0. \quad (8)$$

¹We have the relations $E^i = F^{0i} = -\frac{1}{2} \varepsilon^{ijk} \tilde{F}^{jk}$, and $B^i = \tilde{F}^{0i} = \frac{1}{2} \varepsilon^{ijk} F^{jk}$. I.e. the duality transformation $F^{\mu\nu} \rightarrow \tilde{F}^{\mu\nu}$ amounts to $(\mathbf{E}, \mathbf{B}) \rightarrow (\mathbf{B}, -\mathbf{E})$.

g) The Lagrangian density \mathcal{L} is invariant under the transformation

$$\mathbf{A}(\mathbf{x}, t) \rightarrow \mathbf{A}'(\mathbf{x}, t) = \mathbf{A}(\mathbf{x}, t) + \nabla \Lambda(\mathbf{x}),$$

for all differentiable functions $\Lambda(\mathbf{x})$. Use the Nöther theorem to find the corresponding conserved Nöther current J_Λ .

The general expression for the Nöther current is

$$J^\mu = \frac{\partial \mathcal{L}}{\partial \partial_\mu \Phi_a} \delta \Phi_a, \quad (9)$$

where Φ_a runs over all available fields. Here we have $\delta \varphi = 0$ and $\delta \mathbf{A} = \nabla \Lambda$. Thus we find

$$J_\Lambda^0 = \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{A}}} \cdot \nabla \Lambda = -\lambda \varphi \mathbf{B} \cdot \nabla \Lambda, \quad (10)$$

$$J_\Lambda^k = \lambda \varphi \varepsilon^{jk\ell} E^j \partial_\ell \Lambda = -\lambda \varphi (\mathbf{E} \times \nabla \Lambda)^k. \quad (11)$$

Comment: By introducing the dual field tensor this can also be written in manifestly covariant form

$$J_\Lambda^\mu = -\lambda \varphi \tilde{F}^{\mu\nu} \partial_\nu \Lambda. \quad (12)$$

Problem 2.

The field expansion of the free electromagnetic field in Coulomb gauge is

$$\mathbf{A}(x) = \sum_{\mathbf{k}, r} \frac{1}{\sqrt{2|\mathbf{k}|V}} \left(a_{\mathbf{k}, r} \hat{e}_{\mathbf{k}, r} e^{-ikx} + \text{hermitian conjugate} \right). \quad (13)$$

Then the matrix element $\langle \Omega | a_{\mathbf{q}, s} \mathbf{A}(x) | \Omega \rangle$ equals

- A. 0 ☐
- B. $\frac{1}{\sqrt{2|\mathbf{q}|V}} \hat{e}_{\mathbf{q}, s} e^{-iqx}$ ☐
- C. $a_{\mathbf{q}, s}$ ☐
- D. $\frac{1}{\sqrt{2|\mathbf{q}|V}} \hat{e}_{\mathbf{q}, s}^* e^{iqx}$ ☒
- E. None of the alternatives above. ☐

Problem 3.

Let \mathcal{T} be the time ordering operator, and $\varphi(x)$, $\varphi^\dagger(x)$ quantized complex Klein Gordon fields. Then we have (in natural units, i.e. when $\hbar = c = 1$)

- A. $\mathcal{T} \{ \varphi(x) \varphi^\dagger(y) \} = \mathcal{T} \{ \varphi^\dagger(y) \varphi(x) \}$ ☒
- B. $\mathcal{T} \{ \varphi(x) \varphi^\dagger(y) \} = \mathcal{T} \{ \varphi^\dagger(y) \varphi(x) \} + iG_F(x - y)$ ☐
- C. $\mathcal{T} \{ \varphi(x) \varphi^\dagger(y) \} = \mathcal{T} \{ \varphi^\dagger(y) \varphi(x) \} - iG_F(x - y)$ ☐
- D. $\mathcal{T} \{ \varphi(x) \varphi^\dagger(y) \} = -\mathcal{T} \{ \varphi^\dagger(y) \varphi(x) \} - iG_F(x - y)$ ☐
- E. None of the alternatives above. ☐

Here $G_F(x - y)$ is the Feynman propagator for a complex Klein Gordon field.

Problem 4.

The Dirac equation

$$[i(\gamma^0 \partial_0 + \boldsymbol{\gamma} \cdot \boldsymbol{\nabla}) - m] \psi(x^0, \mathbf{x}) = 0$$

is invariant under space inversion (parity transformation), $\mathbf{x} \rightarrow -\mathbf{x}$. I.e, if $\psi(x^0, \mathbf{x})$ solves the Dirac equation then so does $\psi_P(x^0, \mathbf{x})$, where

- A. $\psi_P(x^0, \mathbf{x}) = i\gamma^2 \psi^*(x^0, -\mathbf{x})$ ☐
- B. $\psi_P(x^0, \mathbf{x}) = \gamma^1 \gamma^3 \psi^*(x^0, -\mathbf{x})$ ☐
- C. $\psi_P(x^0, \mathbf{x}) = \psi(x^0, -\mathbf{x})$ ☐
- D. $\psi_P(x^0, \mathbf{x}) = \gamma^0 \psi(x^0, -\mathbf{x})$ ☒
- E. $\psi_P(x^0, \mathbf{x}) = \psi^*(-x^0, -\mathbf{x})$ ☐