

NTNU Trondheim, Institutt for fysikk**Examination for FY3464 Quantum Field Theory I**

Contact: Jens O. Andersen, tel. 7359 3131

Allowed tools: mathematical tables

1. Noether's theorem.

Show that a continuous global symmetry of a set of fields ϕ_a described by a Lagrangian $\mathcal{L}(\phi_a, \partial_\mu \phi_a)$ leads classically to the conserved current

$$j^\mu = \frac{\delta \mathcal{L}}{\delta \partial_\mu \phi_a} \delta \phi_a - K^\mu, \quad (1)$$

where K^μ is a four-divergence, $\delta \mathcal{L} = \partial_\mu K^\mu$. (6 pts)

We assume first that $\mathcal{L}(\phi_a, \partial_\mu \phi_a)$ is invariant under an infinitesimal change $\delta \phi_a$,

$$0 = \delta \mathcal{L} = \frac{\delta \mathcal{L}}{\delta \phi_a} \delta \phi_a + \frac{\delta \mathcal{L}}{\delta \partial_\mu \phi_a} \delta \partial_\mu \phi_a. \quad (2)$$

Exchange $\delta \partial_\mu = \partial_\mu \delta$ in the second term and use then the Lagrange equations, $\delta \mathcal{L} / \delta \phi_a = \partial_\mu (\delta \mathcal{L} / \delta \partial_\mu \phi_a)$, in the first one. Combining the two terms using the product rule gives

$$0 = \delta \mathcal{L} = \partial_\mu \left(\frac{\delta \mathcal{L}}{\delta \partial_\mu \phi_a} \right) \delta \phi_a + \frac{\delta \mathcal{L}}{\delta \partial_\mu \phi_a} \partial_\mu \delta \phi_a = \partial_\mu \left(\frac{\delta \mathcal{L}}{\delta \partial_\mu \phi_a} \delta \phi_a \right). \quad (3)$$

Hence the invariance of \mathcal{L} under the change $\delta \phi_a$ implies the existence of a conserved current, $\partial_\mu j^\mu = 0$, with

$$j^\mu = \frac{\delta \mathcal{L}}{\delta \partial_\mu \phi_a} \delta \phi_a. \quad (4)$$

If the transformation $\delta \phi_a$ leads to change in \mathcal{L} that is a total four-divergence, $\delta \mathcal{L} = \partial_\mu K^\mu$, then the equations of motion remain invariant. The conserved current j^μ is changed to

$$j^\mu = \frac{\delta \mathcal{L}}{\delta \partial_\mu \phi_a} \delta \phi_a - K^\mu. \quad (5)$$

2. A complex scalar field.

Consider a complex, scalar field ϕ with mass m and a quartic self-interaction proportional to λ in $d = 4$ space-time dimensions.

a.) Write down its Lagrange density \mathcal{L}_s , explain your choice of signs and pre-factors (when physically relevant). (6 pts)

- b.) Determine the mass dimension of all quantities in the Lagrange density \mathcal{L}_s . (6 pts)
 c.) Show that the Lagrange density \mathcal{L}_s is invariant under global phase transformations and determine the conserved current j^μ . (6 pts)

a.) It is easiest to start from two real fields ϕ_1 and ϕ_2 , combining them afterwards into the complex field $\phi = (\phi_1 + i\phi_2)/\sqrt{2}$.

We have to decide which signature we use for the metric, and choose $(+, -, -, -)$. A Lagrange function has the form $L = T - V$, and thus $\dot{\phi}^2$ should have a positive coefficient, while all other terms are negative. Thus we choose the Lagrange density as

$$\mathcal{L}_i = A(\dot{\phi}_i^2 - (\nabla\phi_i)^2) - Bm^2\phi_i^2 - C\frac{\lambda}{4!}\phi_i^4 \quad (6)$$

with $A, B, C > 0$ (Lorentz invariance requires that the coefficient of $\dot{\phi}^2$ and $(\nabla\phi)^2$ agree). This choice of signs can be confirmed by calculating the Hamiltonian density \mathcal{H} , and requiring that it is bounded from below and stable against small perturbations. The correct dispersion relation for a free particle requires $A = B$. The kinetic energy of a canonically normalised field has the coefficient $A = 1/2$; this gives the correct size of vacuum fluctuations. The choice of $C > 0$ is arbitrary. Expressed by the complex fields, the Lagrangian becomes

$$\mathcal{L}_s = \partial_\mu\phi^\dagger\partial^\mu\phi - m^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2. \quad (7)$$

(Alternatively, we could have used that unitarity requires that \mathcal{L} is real. Thus we should use bilinear quantities $\phi^*\phi$ or $\partial_\mu\phi^*\partial^\mu\phi$.)

b.) The action $S = \int dx\mathcal{L}$ enters as $\exp(i/\hbar S)$ the path integral and is therefore in natural units dimensionless. Thus \mathcal{L} has mass dimension 4. From the kinetic term, we see that the mass dimension of the scalar field is 1. Thus the mass dimension of m is, not surprisingly, 1, and λ is dimensionless.

c.) Under the combined global phase transformations $\phi \rightarrow e^{i\vartheta}\phi$ and $\phi^\dagger \rightarrow e^{-i\vartheta}\phi^\dagger$, the Lagrangian \mathcal{L}_s is clearly invariant. With $\delta\phi = i\phi$, $\delta\phi^\dagger = -i\phi^\dagger$, the conserved current follows as

$$j^\mu \propto i \left[\phi^\dagger \partial^\mu \phi - (\partial^\mu \phi^\dagger) \phi \right]. \quad (8)$$

[We can drop the real parameter ϑ , but should keep the imaginary unit i such that the charge is real.]

3. Scalar QED.

Consider now the complex, scalar field ϕ coupled to the photon A^μ , i.e. a massless spin-1 field which field-strength satisfies $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$.

- a.) Find the coupling \mathcal{L}_I between ϕ and A^μ requiring that $\mathcal{L}_s + \mathcal{L}_I$ is invariant under local phase transformations; determine the transformation law for A_μ . (10 pts)
 b.) Write down the generating functionals for disconnected and connected Green functions of this theory. (6 pts)

- c.) How does one obtain connected Green functions for the photon from the generating functional? (4 pts)
- d.) Find the Feynman rules for the vertices involving photons and scalars. (6 pts)
- e.) Define the superficial degree of divergence D and draw for each of the cases $D = \{0, 1, 2\}$ one 1-loop Feynman diagram. (7 pts)

a.) In case of global transformations, $\phi(x) \rightarrow \psi'(x) = U\phi(x)$, the normal derivative transformed as $\partial_\mu \phi(x) \rightarrow [\partial_\mu \phi(x)]' = U[\partial_\mu \psi(x)]$ and thus the kinetic energy was invariant. We define a new covariant derivative D_μ requiring

$$D_\mu \phi(x) \rightarrow [D_\mu \phi(x)]' = U(x)[D_\mu \phi(x)] \quad (9)$$

such that $(D_\mu \phi')^\dagger (D^\mu \phi') = (D_\mu \phi)^\dagger U^\dagger U (D^\mu \phi) = (D_\mu \phi)^\dagger D^\mu \phi$. The gauge field should compensate the difference between the normal and the covariant derivative,

$$D_\mu \phi(x) = [\partial_\mu + iqA_\mu(x)]\phi(x). \quad (10)$$

Now we determine the transformation properties of D_μ and A_μ demanding that $\phi'(x) = U(x)\phi(x)$ and (9) hold. Combining both requirements gives

$$D_\mu \phi(x) \rightarrow [D_\mu \phi]' = U D_\mu \phi = U D_\mu U^{-1} U \phi = U D_\mu U^{-1} \phi', \quad (11)$$

and thus the covariant derivative transforms as $D'_\mu = U D_\mu U^{-1}$. Using its definition (10), we find

$$[D_\mu \phi]' = [\partial_\mu + iqA'_\mu]U\phi = U D_\mu \phi = U[\partial_\mu + iqA_\mu]\phi. \quad (12)$$

[Although not necessary, we do not use that A_μ is abelian.] Compare the second and the fourth term, after having performed the differentiation $\partial_\mu(U\phi)$. The result

$$[(\partial_\mu U) + iqA'_\mu U]\phi = iqU A_\mu \phi \quad (13)$$

should be valid for arbitrary ϕ and hence we arrive after multiplying from the right with U^\dagger at

$$A_\mu \rightarrow A'_\mu = U A_\mu U^\dagger + \frac{i}{g}(\partial_\mu U)U^\dagger = A_\mu - \partial_\mu \vartheta$$

where the last step is valid for an abelian transformation as in the case of a photon.

We obtain \mathcal{L}_I by multiplying out the covariant derivatives,

$$(D_\mu \phi)^\dagger D^\mu \phi = (\partial_\mu \phi)^\dagger \partial^\mu \phi - iq \left[\phi^\dagger \partial^\mu \phi - (\partial^\mu \phi^\dagger) \phi \right] A_\mu + q^2 \phi^\dagger \phi A_\mu A^\mu = (\partial_\mu \phi)^\dagger \partial^\mu \phi + \mathcal{L}_I.$$

[If you use as start the ansatz $\mathcal{L}_I = qA_\mu j^\mu$, you have to realise that the replacement $\partial_\mu \rightarrow D_\mu$ is necessary in the Noether current.]

b.) The complete classical Lagrangian is

$$\mathcal{L}_{\text{cl}} = \mathcal{L}_s + \mathcal{L}_I - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}.$$

To make the path-integral in a covariant gauge well-defined for an abelian gauge field, it is sufficient to add a gauge-fixing term

$$\mathcal{L}_{\text{gf}} = -\frac{1}{2\xi}(\partial^\mu A_\mu)^2.$$

Finally, we add to \mathcal{L}_{eff} sources J_μ coupled linearly to the fields,

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{cl}} + \mathcal{L}_{\text{gf}} + J^\dagger \phi + \phi^\dagger J + J^\mu A_\mu. \quad (14)$$

The generating functional Z for disconnected Green functions is the path integral over fields over $\exp(i \int d^4x \mathcal{L}_{\text{eff}})$,

$$Z[J, J^*, J^\mu] = \int \mathcal{D}\phi \mathcal{D}\phi^* \mathcal{D}A_\mu \exp\{i \int d^4x \mathcal{L}_{\text{eff}}\} = e^{iW[J, J^*, J^\mu]}. \quad (15)$$

[We assume implicitly the Feynman prescription to ensure the convergence of the path integral.]

c.) $W[J, J^*, J^\mu]$ generates connected Green functions for the photon via

$$\frac{1}{i^n} \frac{\delta^n W}{\delta J_\mu(x_1) \cdots \delta J_\nu(x_n)} \Big|_{J=0} = G_{\mu \cdots \nu}(x_1, \dots, x_n). \quad (16)$$

d.) We go to momentum space performing a Fourier transformation of \mathcal{L}_I . For the trilinear vertex,

$$\begin{aligned} F &= -iq \int d^4p_1 d^4p_2 d^4p_3 (2\pi)^4 \delta(p_1 - p_2 - p_3) \left[\phi^\dagger(p_1) \partial^\mu \phi(p_2) - (\partial^\mu \phi^\dagger(p_1) \phi(p_2)) \right] A_\mu(p_3) \\ &= -iq(-ip_2^\mu - ip_1^\mu) \phi^\dagger(p_1) \phi(p_2) A_\mu(p_3) \end{aligned} \quad (17)$$

Multiplying by i and taken derivatives w.r.t. to the three fields, we can read off as rule for the $\phi^\dagger \phi A_\mu$ vertex in case of 2 ϕ particles $-iq(p_1^\mu + p_2^\mu)$. Since antiparticles propagate backwards, their momenta enters with a minus sign. Proceeding similarly for the quartic vertex, $\phi^\dagger \phi A_\mu A^\mu$, we find $2iq^2 \eta_{\mu\nu}$.

e.) The superficial degree D of divergence of a 1PI Feynman graph is given by the number of momenta in the numerator minus the number of momenta in the denominator. In $d = 4$ space-time dimensions, L independent loop momenta contribute thus $4L$ and derivative couplings a factor d_V counting the power of derivatives from all vertices, while I internal bosonic lines subtract the factor $2I$. Combined



$$D = 4L + d_V - 2I.$$

where L is the number of independent loop momenta and I the number of internal lines. Examples are:


- $D = 0$



- $D = 1$



- $D = 2$



4. Tensor decomposition.

Consider the *real* decay process $\mu \rightarrow e + \gamma$ in Minkowski space, allowing for parity violation. Write its Lorentz invariant amplitude \mathcal{A} as $\mathcal{A}(\mu \rightarrow e + \gamma) = \varepsilon_\lambda \langle e | J_{\text{em}}^\lambda | \mu \rangle$ where J_{em}^λ denotes the electromagnetic current and decompose it in scalar functions A, B, \dots as

$$\langle e | J_{\text{em}}^\lambda | \mu \rangle = \bar{u}_e(p') [A \gamma^\lambda + \dots] u_\mu(p).$$

Use the symmetries to express \mathcal{A} by the minimal number of scalar functions required. [Note the difference to the treatment of the electromagnetic vertex in the lectures where the photon was virtual.] (10 pts)

Translation invariance of Minkowski space implies four-momentum conservation, i.e. $q = p' - p$. Translation invariance combined with the on-shell condition means that the photon sees only the momentum difference $q = p' - p$, not the individual momenta p and p' . Hence J_{em}^λ can be only a function of the momentum difference, $J_{\text{em}}^\lambda = J_{\text{em}}^\lambda(q^\mu)$, while the arbitrary scalar functions can depend only on q^2 .

We have to form all possible vectors out of the momenta q_μ and the 16 basis elements

$$\Gamma_i = \{1, \gamma^\mu, \gamma^\mu \gamma^5, \sigma^{\mu\nu}, \gamma^5\}$$

of the Clifford algebra.

$$J_{\text{em}}^\mu(q) = (A + B \gamma^5) \gamma^\mu + (C + D \gamma^5) q^\mu + (D + E \gamma^5) i \sigma^{\mu\nu} q_\nu.$$

The electromagnetic current J_{em}^λ is conserved, $\partial_\lambda J_{\text{em}}^\lambda(x) = 0$ or $q_\lambda J_{\text{em}}^\lambda(q) = 0$. This gives the condition

$$-m_e(A + B\gamma^5) + m_\mu(A - B\gamma^5) + q^2(C + D\gamma^5) = 0$$

or

$$A = B = 0$$

as the photon is on-shell, $q^2 = 0$. Finally we use that the photon is transverse, $\varepsilon^\mu q_\mu = 0$, to obtain

$$\mathcal{A}(\mu \rightarrow e + \gamma) = \varepsilon_\lambda \langle e | J_{\text{em}}^\lambda | \mu \rangle = \varepsilon_\lambda \bar{u}_e(p') [(D + E\gamma^5) i\sigma^{\lambda\nu} q_\nu] u_\mu(p).$$

Thus only the on-shell values of the magnetic form factors, $D(q^2 = 0)$ and $E(q^2 = 0)$, contribute to the decay process.

Some formulas

The Gamma matrices satisfy the Clifford algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} \quad (18)$$

and are in the Weyl or chiral representation given by

$$\gamma^0 = 1 \otimes \tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (19)$$

$$\gamma^i = \sigma^i \otimes i\tau_3 = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}, \quad (20)$$

$$\gamma^5 = 1 \otimes \tau_3 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (21)$$

$$\psi_L = \frac{1}{2}(1 - \gamma^5)\psi \equiv P_L\psi \quad \text{and} \quad \psi_R = \frac{1}{2}(1 + \gamma^5)\psi \equiv P_R\psi. \quad (22)$$

$$\sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu] \quad (23)$$