

NTNU Trondheim, Institutt for fysikk

Examination for FY3464 Quantum Field Theory I

Contact: Michael Kachelrieß, tel. 99890701

Allowed tools: mathematical tables

1. Scalar field and scale invariance.

Consider a complex, scalar field ϕ with mass m and self-interaction $g\phi^n$.

- Write down the Lagrange density \mathcal{L} , explain your choice of signs and pre-factors (when physically relevant). (5 pts)
- Determine the mass dimension in $d = 4$ space-time dimensions of all quantities in the Lagrange density \mathcal{L} . Choose n such that the coupling g is dimensionless. (5 pts)
- Set now $m = 0$ and consider a real scalar field ϕ . Find the equation of motion for $\phi(x)$. (4 pts)
- Assume that $\phi(x)$ solves the equation of motion and define a scaled field

$$\tilde{\phi}(x) \equiv e^{D^a} \phi(e^a x), \quad (1)$$

where D and a are constants. Show that the scaled field $\tilde{\phi}(x)$ is also a solution of the equation of motion, provided that the constant D is chosen appropriately. (6 pts)

- Bonus question: Argue, if the classical symmetry (1) is (not) conserved on the quantum level. [max. 50 words] (2 pts)

- The free Lagrangian of a real scalar field is

$$\mathcal{L}_0 = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2.$$

The relative sign is fixed by the relativistic energy-momentum relation, the overall sign by the requirement that the Hamiltonian is bounded from below. The factor $1/2$ in the kinetic energy leads to “canonically normalised” field, The factor $1/2$ for the mass follows then from the relativistic energy-momentum relation. Combining two real fields into a complex one, $\phi = (\phi_1 + i\phi_2)/\sqrt{2}$ gives

$$\mathcal{L}_0 = \partial_\mu \phi^\dagger \partial^\mu \phi - m^2 \phi^\dagger \phi.$$

If the self-interaction is odd, both choices of sign will lead to an unstable vacuum. If the self-interaction is even, the choice $-g(\phi^\dagger \phi)^{n/2}$ will lead to positive potential energy and thus stability of the vacuum.

- The action $S = \int d^4x \mathcal{L}$ has to be dimensionless. Thus $[\mathcal{L}] = m^4$, $[\partial_\mu \phi] = m^2$, and thus $[\phi] = m$. Thence $[m] = m$, and the coupling is dimensionless, $[\lambda] = m^0$ for $n = 4$.

- Using the Lagrange equation or varying directly the action gives

$$\square \phi + \frac{\lambda}{3!} \phi^3 = 0.$$

d.) Set $y = e^a x$. Then

$$\frac{\partial}{\partial x^\mu} = \frac{\partial y^\mu}{\partial x^\mu} \frac{\partial}{\partial y^\mu} = e^a \frac{\partial}{\partial y^\mu}$$

and $\square_x = e^{2a} \square_y$. Then $\tilde{\phi}$ satisfies the equation of motion,

$$\square_x \tilde{\phi} + \frac{\lambda}{3!} \tilde{\phi}^3 = e^{(2+D)a} \square_x \phi + e^{3Da} \frac{\lambda}{3!} \phi^3 \stackrel{!}{=} e^{3a} \left[\square_x \phi + \frac{\lambda}{3!} \phi^3 \right] \stackrel{!}{=} 0.$$

if we choose $D = 1$. Thus the scalar field should scale as its “naive” dimension suggests.

e.) Bonus: We discussed in Exercise sheet 7 scale invariance and noted as requirement that the classical Lagrangian contains no dimension-full parameters (which would fix scales). But loop corrections introduce necessarily a scale (μ in DR, Λ as cutoff). As a consequence, scale invariance is broken by quantum corrections.

Remarks: 1. As alternative in b), one can check the transformation of the action; surprisingly, you find then the constraint $D = 1$ and $d = 4$.

2. If we do not assume $a = \text{const.}$, we leave Minkowski space and have to consider the scalar field in a general space-time. Then one finds that the action is invariant under this transformation with an arbitrary, positive function $a(x)$, if one adds (in $d = 4$) a coupling $-R\phi^2/6$ between ϕ and the curvature scalar R .

2. Fermion field.

Consider a massless Dirac field ψ with Lagrangian

$$\mathcal{L} = \bar{\psi}(i\cancel{D})\psi.$$

a.) Derive the propagator $S_F(p)$ of the field ψ . [You do not have to discuss how the poles of $S_F(p)$ are treated.] (4 pts)

b.) Write down the generating functional for disconnected Green functions for this theory. (4 pts)

c.) Show that the Lagrange density \mathcal{L} is invariant under global vector phase transformations $U_V(1)$, $\psi \rightarrow \psi' = e^{i\theta}\psi$, and under global axial phase transformations $U_A(1)$, $\psi \rightarrow \psi' = e^{i\theta\gamma^5}\psi$. (6 pts)

d.) Show that global symmetry under vector phase transformation $U_V(1)$ can be made local, if a coupling to a gauge boson is added. (5 pts)

e.) Draw the divergent one-loop diagrams and determine their superficial degree of divergence D (in $d = 4$ space-time dimensions) for the theory coupled to a gauge boson. (8 pts)

a.) The Green functions of the free Dirac equation (for $m > 0$) are defined by

$$(i\cancel{D} - m)S(x, x') = \delta(x - x'), \quad (2)$$

where we omit on the RHS a unit matrix in spinor space. Translation invariance implies $S(x, x') = S(x - x')$ and, performing a Fourier transformation, the Fourier components $S(p)$ have to obey

$$(\not{p} - m)S(p) = 1. \quad (3)$$

After multiplication with $\not{p} + m$ and use of $\not{p}^2 = \frac{1}{2}\{\gamma^\mu, \gamma^\nu\}a_\mu a_\nu = a^2$, we can solve for the propagator in momentum space, After multiplication with $\not{p} + m$ and use of $\not{p}^2 = \frac{1}{2}\{\gamma^\mu, \gamma^\nu\}a_\mu a_\nu = a^2$, we can solve for the propagator in momentum space,

$$iS_F(p) = i \frac{\not{p} + m}{p^2 - m^2 + i\varepsilon} = \frac{i}{\not{p} - m + i\varepsilon}, \quad (4)$$

where the last step is only meant as a symbolical shortcut.

b.) The path integral in phase space is for zero sources

$$Z[0] = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{iS[\psi, \bar{\psi}]} = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{i \int d^4x \bar{\psi} (i\not{\partial} - m) \psi}. \quad (5)$$

Adding Grassmannian sources η and $\bar{\eta}$ gives

$$\begin{aligned} Z[\eta, \bar{\eta}] &= \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{i \int d^4x d^4x' [\bar{\psi}(x') A(x', x) \psi(x) + \bar{\eta}(x') \psi(x) + \bar{\psi}(x') \eta(x)]} = \\ &= Z[0] \exp \left(-i \int d^4x d^4x' \bar{\eta}(x) S_F(x - x') \eta(x') \right). \end{aligned} \quad (6)$$

c.) For vector transformations, $\psi \rightarrow \psi' = e^{i\theta} \psi$ implies $\bar{\psi} \rightarrow \bar{\psi}' = e^{i\theta} \bar{\psi}$ and the global phases drop out. For axial transformations, it follows with $\{\gamma^5, \gamma^\mu\} = 0$,

$$\psi'(x) \rightarrow e^{i\phi\gamma^5} \psi(x) \quad \text{and} \quad \bar{\psi}(x) \rightarrow \bar{\psi}'(x) = (e^{i\phi\gamma^5} \psi(x))^\dagger \gamma^0 = \bar{\psi}(x) e^{i\phi\gamma^5}. \quad (7)$$

Thus \mathcal{L} without mass term is invariant too.

d.) Introduce a covariant derivative,

$$\partial_\mu \rightarrow D_\mu = \partial_\mu + iqA_\mu, \quad (8)$$

which transforms homogeneously,

$$D_\mu \psi(x) \rightarrow D'_\mu \psi'(x) = \{\partial_\mu + iq[A_\mu(x) - \partial_\mu \Lambda(x)]\} \exp[iq\Lambda(x)] \psi(x) = \quad (9)$$

$$= \exp[iq\Lambda(x)] \{\partial_\mu + iqA_\mu(x)\} \psi(x) = U(x) D_\mu \psi(x). \quad (10)$$

e.) The theory corresponds to QED; see e.g. Fig. 11.3 of the notes.

3. Unitarity.

a.) Derive the optical theorem

$$2\Im T_{ii} = \sum_n T_{in}^* T_{ni}.$$

Give a physical interpretation of this relation (less than 50 words).

(6 pts)

b.) The vacuum polarisation of a photon,

$$q \quad \text{---} \circlearrowleft \text{---} q = \Pi^{\mu\nu}(q^2) = (q^2 \eta^{\mu\nu} - q^\mu q^\nu) \Pi(q^2)$$

is given in dimensional regularisation by

$$\Pi(q^2) = -\frac{e^2}{12\pi^2} \left\{ \frac{1}{\varepsilon} - \gamma + \ln(4\pi) - 6 \int_0^1 dx \, x(1-x) \ln \left[\frac{m^2 - q^2 x(1-x)}{\mu^2} \right] \right\}.$$

Show that gauge invariance, $q_\mu \Pi^{\mu\nu}(q) = 0$, implies as tensor structure of the vacuum polarisation tensor $\Pi^{\mu\nu}(q) = (q^2 \eta^{\mu\nu} - q^\mu q^\nu) \Pi(q^2)$. (4 pts)

c.) Derive the imaginary part of the vacuum polarisation, $\Im[\Pi(q^2)]$. (6 pts)

d.) How does the imaginary part of the vacuum polarisation changes, if the renormalisation scheme is changed? (4 pts)

a.) The unitarity of the scattering operator, $S^\dagger S = S S^\dagger = 1$, expresses the fact that we (should) use a complete set of states for the initial and final states in a scattering process,

$$1 = \sum_n |n, +\infty\rangle \langle n, +\infty| = \sum_n S |n, -\infty\rangle \langle n, -\infty| S^\dagger = S S^\dagger. \quad (11)$$

We split the scattering operator S into a diagonal part and the transition operator T , $S = 1 + iT$, and thus

$$1 = (1 + iT)(1 - iT^\dagger) = 1 + i(T - T^\dagger) + TT^\dagger \quad (12)$$

or

$$\mathbf{i}TT^\dagger = T - T^\dagger. \quad (13)$$

We now consider matrix elements between the initial and final state,

$$\langle f|T - T^\dagger|i\rangle = T_{fi} - T_{if}^* = i\langle f|TT^\dagger|i\rangle = i\sum_n T_{fn}T_{in}^*. \quad (14)$$

If we set $|i\rangle = |f\rangle$, we obtain optical theorem as a connection between the forward scattering amplitude T_{ii} and the scattering into all possible states n ,

$$2\Im T_{ii} = \sum_n |T_{in}|^2. \quad (15)$$

It relates the attenuation of a beam of particles in the state i , $dN_i \propto -|\mathcal{T}_{ii}|^2 N_i$, to the probability that they scatter into all possible states n : what is lost, should show up somewhere.

b.) Using the tensor method, we can express $\Pi^{\mu\nu}(q^2)$ as a linear combination of $\eta^{\mu\nu}$ and $q^\mu q^\nu$. Imposing then current conservation fixes the relative sign.

c.) The only relevant part is the term with the log,

$$\Pi(q^2) = \frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \ln \left[1 - \frac{q^2}{m^2} x(1-x) \right] + \dots$$

The simplest option is to find the x range for which the log is negative for a given q^2 , and to use then $\Im \ln(x + i\varepsilon) = -\pi$.

From $1 - \frac{q^2}{m^2}x(1-x) = 0$, it follows $x_{1/2} = \frac{1}{2} \pm \frac{1}{2}\beta$ with $\beta = \sqrt{1 - 4m^2/q^2}$. Then

$$\Im \Pi^{\text{on}}(q^2 + i\varepsilon) = \frac{2\alpha}{\pi}(-\pi) \int_{\frac{1}{2}-\frac{1}{2}\beta}^{\frac{1}{2}+\frac{1}{2}\beta} dx x(1-x) = -\frac{\alpha}{3}\beta(1 + 2m^2/q^2). \quad (16)$$

c.) All the dependence on the renormalisation scheme is contained in local polynomials of the fields and their derivatives. Non-analytic functions like the log term generating the imaginary part are therefore independent of the renormalisation scheme.

Useful formulas

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}. \quad (17)$$

$$\{\gamma^\mu, \gamma^5\} = 0 \quad \text{and} \quad (\gamma^5)^2 = 1. \quad (18)$$

$$\sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu] \quad (19)$$

$$\bar{\Gamma} = \gamma^0 \Gamma^\dagger \gamma^0 \quad (20)$$

$$\frac{1}{ab} = \int_0^1 \frac{dz}{[az + b(1-z)]^2}. \quad (21)$$

$$\frac{1}{k^2 + m^2} = \int_0^\infty ds e^{-s(k^2 + m^2)} \quad (22)$$

$$I_0(\omega, \alpha) = \int \frac{d^{2\omega} k}{(2\pi)^{2\omega}} \frac{1}{[k^2 - m^2 + i\varepsilon]^\alpha} = i \frac{(-1)^\alpha}{(4\pi)^\omega} \frac{\Gamma(\alpha - \omega)}{\Gamma(\alpha)} [m^2 - i\varepsilon]^{\omega - \alpha}. \quad (23)$$

$$I(\omega, 2) = i \frac{1}{(4\pi)^\omega} \frac{\Gamma(2 - \omega)}{\Gamma(2)} \frac{1}{[m^2 - q^2 z(1-z)]^{2-\omega}}. \quad (24)$$

$$\Im \ln(x + i\varepsilon) = -\pi \quad (25)$$