

### Problem 1. (10 points)

#### A. Constraints

What are holonomic constraints? Under what circumstances can we use the Lagrangian formalism for solving a mechanical problem with non-holonomic constraints? What implications does this have in terms of the number of generalized coordinates? Provide a proper written explanation where you may use some equations also. [6 p]

#### B. Total time-derivative of kinetic energy for a falling particle

Consider the following Hamiltonian of a falling particle:  $H = T + V = \frac{p^2}{2m} + mgy$

Derive the total time-derivative of kinetic energy  $T$  by using the Poisson bracket formalism. The vertical coordinate of the particle (mass  $m$ ) is denoted with  $y$  and the related momentum is  $p$ .

Confirm your result by considering the time-derivative of total energy. [4p]

### Problem 2. Sliding particle with air resistance (8 points)

Let us consider a body (mass  $m$ ) sliding down an inclined plane without any friction from the support. The body experiences a drag force that is directly proportional to velocity and has the form  $F = -kmv$ , where  $k$  is a constant. The inclination angle of the plane is  $\theta$ .

(a) Write out the Lagrangian and solve the equation of motion. [3p]

(b) The body starts from rest ( $v_0 = 0$ ). Solve the equation for velocity as a function of time. What is the upper limit (called terminal velocity)? [3p]

**Hint:** Express  $\ddot{x} = \frac{dv}{dt}$  and carry out the integrations.

(c) Let us assume that the body has achieved 90% of its maximum velocity. When does this occur ( $t$ ) and how far ( $x$ ) has the body slid along the plane? [2p]

### Problem 3. Heavy spinning top (8 points)

Let us consider a heavy spinning top as described in the lectures which rotates frictionless around a fixed point on the ground and experiences gravity. The kinetic energy is of the form

$$T = \frac{1}{2}I(\omega_1^2 + \omega_2^2) + \frac{1}{2}I_3\omega_3^2$$

Let us choose the Euler angles  $(\varphi, \theta, \psi)$  as the generalized coordinates. What physical quantities do they represent? The distance of the center of mass from the fixed tip is denoted with  $h$ . Write out the explicit Lagrangian in terms of the generalized coordinates. Which component of  $\vec{\omega}$  is a constant and what does it represent? [4 p]

Derive the Lagrangian equations of motion and the canonical momenta. What are the conserved quantities? [4p]

**Hint:** There is no need to expand the final time-derivative explicitly for a constant quantity. Instead you may use  $\frac{d}{dt}(\dots) = 0$ .

### Problem 4. Special theory of relativity – creation of a kaon particle (8 points)

(a) Starting from the 4-vector for position (event)  $x_\mu$ , derive the momentum 4-vector  $P_\mu$  where its 4<sup>th</sup> component (according to the covariant 3+1 formulation) is expressed in

terms of total energy. Calculate also the Lorentz invariant property  $P_\mu P_\mu$  by assuming rest frame. These will be useful in the following. [3p]

(b) Let us consider a relativistic collision (reaction):  $\pi^+ + n \rightarrow K^+ + \Lambda_0$ .


What is the threshold kinetic energy of the pion ( $\pi^+$ ) to create kaon ( $K^+$ ) at an angle of 90 degrees in the rest frame of the neutron ( $n$ )? Express your solution in terms of particle masses. The inherent nature of the particles is not relevant, just consider them as particles with different masses. [5p]

**Hint:** We have conservation of 4-momentum. Correspondingly, modify the momentum equation such that you will have  $(P_\mu^\Lambda)^2 = P_\mu^\Lambda P_\mu^\Lambda$  alone on the left-hand side. Next, consider the individual terms taking into account the above-mentioned conditions.

(Final answer:  $\frac{T_\pi}{c^2} \geq \frac{m_\Lambda^2 - m_\pi^2 - m_n^2 - m_K^2 + 2m_n m_K}{2(m_n - m_K)} - m_\pi$ )

### Problem 5. Coriolis effect from another perspective (12 points)

Let us revisit the problem of dropping a pellet down from a tower as seen from outer space (inertial frame), see the figure. Let us denote that latitude on Earth is  $\lambda$  and angular velocity of rotation is  $\omega$ . The radius of Earth is  $R$  and the height of the tower is  $h$ . The pellet starts falling down from rest (tower) in the rotating non-inertial frame of Earth.



**Note:** Points (a)-(c) are straightforward. Come back to (d)-(f) later if you are short of time.

(a) Consider for a short moment that Earth is represented by a single mass point in its center. What shape does the trajectory of the falling particle have? In reality, the particle is able to travel only a small sequence of this trajectory before hitting the ground ( $r = R$ ). What location does the particle have on this orbit in the beginning (tower)? Write out the horizontal velocity  $v_{hor}$  of the particle and the corresponding angular momentum  $\ell$  in the beginning ( $r = R + h$ ), as seen from outside (inertial frame). [2p]

(b) Going back to the solution of the Kepler problem, choose that the orbit angle is  $\theta = 0$  in the beginning and use

$$\frac{p}{r} = 1 - \varepsilon \cos \theta,$$

Show that the distance  $r$  has the general solution [2p]

$$r = \frac{(1 - \varepsilon)(R + h)}{1 - \varepsilon \cos \theta}$$

(c) Use the constant areal velocity (Kepler II) in order to derive the following expression for time [2p]

$$t = \frac{1}{\omega \cos \lambda} \int_0^\theta \frac{(1 - \varepsilon)^2}{(1 - \varepsilon \cos \theta)^2} d\theta$$

(d) Denote the ground landing point as  $\theta = \theta_0$  ( $r = R$ ). Show that [2p]

$$\frac{R+h}{R} = 1 + \frac{2\varepsilon}{1-\varepsilon} \sin^2 \frac{\theta_0}{2} \Rightarrow \frac{h}{R} \cong \frac{\varepsilon \theta_0^2}{2(1-\varepsilon)}$$

**Hint:** Use the trigonometric relation:  $\cos \theta = 1 - 2\sin^2(\theta/2)$  and a Taylor expansion in the last step ( $\theta_0$  small).

(e) Modify the equation for time in (c) for the actual falling time  $t(\theta = \theta_0) = T$  such that it becomes [2p]

$$T \cong \frac{1}{\omega \cos \lambda} \int_0^{\theta_0} \frac{d\theta}{[1 + (h\theta^2/R\theta_0^2)]^2}$$

(f) We can now take a Taylor expansion, integrate, and solve  $\theta_0$  (+ Taylor once more) such that it becomes (you do not need to show this!)

$$\theta_0 \cong \omega T \cos \lambda \left(1 + \frac{2h}{3R}\right)$$

By using this result, calculate the deflection  $d$  of the landing point on the ground as a function of tower height  $h$ . What is the direction of deflection? [2p]

**Hint:** Free fall without air resistance in a non-rotating frame gives you an expression (approximation) for time  $T$  that you can use here for the final step.

*Fun fact: The result will be identically the same as what we obtained earlier in the exercises by using the expression of Coriolis force in a rotating frame!*

GOOD LUCK!

MERRY CHRISTMAS AND HAPPY NEW YEAR!