

TFY4345 Classical Mechanics Exam November 27, 2023

Part I (2.5 points for each correct answer. Answer Part I in the table in Inspira.)

1.1 The constraint $r = \sqrt{x^2 + y^2}$ for a particle sliding on a ring of radius r is called

- A) conservative B) holonomic C) canonical D) invariant E) cyclic F) virtual

1.2 What is the value of the element ε_{321} of the Levi-Civita tensor?

- A) 1 B) -1 C) i D) $-i$ E) 0 F) π

1.3 What is the SI unit of the conjugate momentum to the polar angle θ ?

- A) The same unit as for angular momentum
B) The same unit as for power
C) The same unit as for acceleration
D) Nm
E) m/s^2
F) kg m/s

1.4 If the system Lagrangian is independent of a coordinate q , this coordinate is called

- A) conservative B) holonomic C) canonical D) invariant E) cyclic F) virtual


1.5 A particle with mass m moves in the xy plane in a potential $V(r) = kr^2/2$. Here, k is a positive constant, $r^2 = x^2 + y^2$, $x = r \cos \theta$ and $y = r \sin \theta$. Which quantity (in addition to the total energy) is conserved for this particle?

- A) p_x B) p_y C) v_x D) v_y E) p_θ F) None

1.6 Three point particles moving in three dimensional space are subject to three independent holonomic constraints. How many independent coordinates q_j are needed to describe this system?


- A) 3 B) 4 C) 5 D) 6 E) 7 F) 8

1.7 A triple planar pendulum consists of three balls (point masses) connected by two massless rods of fixed length, and by a third massless rod to the support at A, which may slide without friction in the x direction. The balls are allowed to move in the xy plane. How many independent coordinates q_j are needed to describe this system?



- A) 3 B) 4 C) 5 D) 6 E) 7 F) 8

1.8 Two balls (point masses m_1 and m_2) are connected with a massless rope of length $\ell + \pi R$ that can slide without friction over a cylinder with radius R . Zero potential is chosen at the centre of the cylinder, a distance x above m_1 . What is the Lagrangian $L = T - V$ for this system?



- A) $L = \frac{1}{2}(m_1 - m_2)\dot{x}^2 + (m_1 - m_2)gx + m_2g\ell$
- B) $L = \frac{1}{2}(m_1 - m_2)\dot{x}^2 + (m_1 + m_2)gx + m_2g\ell$
- C) $L = \frac{1}{2}(m_1 + m_2)\dot{x}^2 + (m_1 - m_2)gx + m_2g\ell$
- D) $L = \frac{1}{2}(m_1 + m_2)\dot{x}^2 + (m_1 + m_2)gx + m_2g\ell$
- E) $L = \frac{1}{2}(m_1 + m_2)\dot{x}^2 + (m_1 - m_2)gx + (m_1 + m_2)g\ell$
- F) $L = \frac{1}{2}(m_1 + m_2)\dot{x}^2 + (m_1 - m_2)gx + (m_1 - m_2)g\ell$

1.9 A particle with mass m and charge q moving in an electromagnetic field has the Lagrangian

$$L = \frac{1}{2}m\dot{x}_j\dot{x}_j + qA_j\dot{x}_j - q\phi.$$

What is the canonical momentum p_2 ?

- A) $p_2 = q\dot{x}_2 + mA_2$
- B) $p_2 = m\dot{x}_1 - qA_3$
- C) $p_2 = \dot{x}_2 - A_2$
- D) $p_2 = 2m\dot{x}_2 + qA_2$
- E) $p_2 = 2m\dot{x}_2 + qA_2/2$
- F) $p_2 = m\dot{x}_2 + qA_2$

1.10 If the vector potential is $\mathbf{A} = B_0(x\hat{y} - y\hat{x})$, what is the magnetic field \mathbf{B} ?

- A) $\mathbf{B} = B_0\hat{z}$
- B) $\mathbf{B} = 2B_0\hat{z}$
- C) $\mathbf{B} = (B_0/2)\hat{z}$
- D) $\mathbf{B} = 4B_0\hat{z}$
- E) $\mathbf{B} = (B_0/4)\hat{z}$
- F) $\mathbf{B} = 0$

1.11 If the vector potential is $\mathbf{A} = E_0t\hat{x}$ and the scalar potential is $\phi = E_0(y+z)$, what is the electric field \mathbf{E} ?

- A) $\mathbf{E} = -E_0\hat{x}$
- B) $\mathbf{E} = -E_0(\hat{y} + \hat{z})$
- C) $\mathbf{E} = -E_0(\hat{x} + \hat{z})$
- D) $\mathbf{E} = -E_0(\hat{x} + \hat{y})$
- E) $\mathbf{E} = -E_0(\hat{x} + \hat{y} + \hat{z})$
- F) $\mathbf{E} = 0$

1.12 If the Lagrangian for a system with two independent coordinates q_1 and q_2 is

$$L = c_1\dot{q}_1^2 + c_2\dot{q}_2^2 + c_3q_1\dot{q}_1 + c_4\dot{q}_2^2,$$

what is the canonical momentum p_1 ?

- A) $p_1 = c_1q_1$
- B) $p_1 = c_2q_1$
- C) $p_1 = c_3q_1$
- D) $p_1 = c_1q_2$
- E) $p_1 = c_2q_2$
- F) $p_1 = c_3q_2$

1.13 A planet moves in an elliptical orbit with eccentricity 0.21 around a much heavier star located at the origin. What is the ratio r_{\min}/r_{\max} between the shortest and longest distance from the planet to the star?

- A) 0.45
- B) 0.55
- C) 0.65
- D) 0.75
- E) 0.85
- F) 0.95

1.14 Which transformation matrix describes rotation an angle ϕ counterclockwise around the x_2 axis?

A) $\begin{pmatrix} \cos \phi & 0 & -\sin \phi \\ 0 & 1 & 0 \\ \sin \phi & 0 & \cos \phi \end{pmatrix}$

B) $\begin{pmatrix} \cos \phi & 0 & \sin \phi \\ 0 & 1 & 0 \\ \sin \phi & 0 & -\cos \phi \end{pmatrix}$

C) $\begin{pmatrix} \cos \phi & 0 & -\sin \phi \\ 0 & 1 & 0 \\ -\sin \phi & 0 & \cos \phi \end{pmatrix}$

D) $\begin{pmatrix} \sin \phi & 0 & -\cos \phi \\ 0 & 1 & 0 \\ \cos \phi & 0 & \sin \phi \end{pmatrix}$

E) $\begin{pmatrix} -\sin \phi & 0 & -\cos \phi \\ 0 & 1 & 0 \\ \cos \phi & 0 & \sin \phi \end{pmatrix}$

F) $\begin{pmatrix} -\cos \phi & 0 & -\sin \phi \\ 0 & 1 & 0 \\ -\sin \phi & 0 & -\cos \phi \end{pmatrix}$

1.15 A spaceship moves with speed $7c/8$ relative to Spaceman Spiff, who is at rest on Anhooie-4. The alien Hideous Blob is shot out of the spaceship, in the forward direction, with speed $5c/6$ relative to the spaceship. What is the speed of Hideous Blob as observed by Spaceman Spiff?

- A) $82c/83$ B) $6c/7$ C) $41c/24$ D) $7c/8$ E) $35c/48$ F) $c/3$

1.16 The acceleration \mathbf{a}_b of an object measured by a stationary observer at the Equator on the surface of the Earth is

$$\mathbf{a}_b = \mathbf{a}_s + 2\mathbf{v}_b \times \boldsymbol{\omega} - \boldsymbol{\omega} \times (\boldsymbol{\omega} \times \mathbf{r}).$$

Here, \mathbf{v}_b and \mathbf{r} are the velocity and the position of the object, respectively, both measured by this observer, and \mathbf{a}_s is the acceleration measured in an inertial system. Assume a spherical Earth with radius $r = 6378$ km rotating around an axis pointing straight north, with a period $T = 2\pi/\omega = 24$ hours. The observer throws a ball upwards with an initial speed 15 m/s. At this instant, what is the Coriolis acceleration?

- A) 5.5 mm/s^2 B) 4.4 mm/s^2 C) 3.3 mm/s^2 D) 2.2 mm/s^2 E) 1.1 mm/s^2 F) 0.55 mm/s^2

1.17 At the instant described in the previous question, what is the centrifugal acceleration?

- A) 64 mm/s^2 B) 54 mm/s^2 C) 44 mm/s^2 D) 34 mm/s^2 E) 24 mm/s^2 F) 14 mm/s^2

1.18 What is the mass of a (free) particle with energy 500 MeV and momentum 400 MeV/c?

- A) 50 MeV/c² B) 100 MeV/c² C) 150 MeV/c²
D) 200 MeV/c² E) 250 MeV/c² F) 300 MeV/c²

1.19 In a canonical transformation of type 1, from "old" coordinates (q, p) to "new" coordinates (Q, P) , the generating function is $F = F_1(q, Q) = qQ - q^2 - Q^2$. What is $P(q, p)$?


- A) $P = q - p$ B) $P = 4q + 3p$ C) $P = q + p$
D) $P = 2q - 2p$ E) $P = 4q - 3p$ F) $P = 3q + 2p$

1.20 What is most likely a key ingredient in Unni's Cinnamon Cake?

- A) Anise
B) Basil
C) Cinnamon
D) Dill
E) Estragon
F) Fennel



Part II (Weights are given for each of the 8 partial questions)



2 (20%) Three balls with masses m , $2m$ and $4m$ ($m = 50$ g) are connected by identical and ideal springs with spring constant $k = 150$ N/m, as shown in the figure above. The balls can move along the x axis only, and we consider small oscillations around their equilibrium positions x_{01} , x_{02} and x_{03} .

a) (7%) With the balls' deviations from equilibrium, $\eta_i = x_i - x_{0i}$, as coordinates, the potential V and kinetic energy T are both quadratic forms,

$$V = \frac{1}{2} V_{ij} \eta_i \eta_j \quad \text{and} \quad T = \frac{1}{2} T_{ij} \dot{\eta}_i \dot{\eta}_j,$$

respectively. Determine the 3×3 symmetric matrix \mathbf{V} and diagonal matrix \mathbf{T} , with matrix elements V_{ij} and T_{ij} .

b) (7%) Solve the secular equation


$$|\mathbf{V} - \omega^2 \mathbf{T}| = 0$$

(i.e., zero determinant) and determine the two nonzero eigenfrequencies $f_j = \omega_j/2\pi$ ($j = 1, 2$) of this system. (Determine both numerical values and units.)

Hint: You will end up with a 3rd order equation for ω^2 , where one root is $\omega^2 = 0$. You may find it convenient to extract a factor k^3 from the determinant and introduce the dimensionless variable $\alpha = m\omega^2/k$.

c) (6%) Determine the amplitudes (i.e., relative amplitudes, including sign) of the three balls in the normal mode with the *smallest* eigenfrequency.

3 (25%) A rod with negligible mass and length ℓ has equal masses m at its two ends. (See figure below.) One mass can slide without friction along a horizontal constraint on the x axis. The other mass is restricted to move in the xy plane. We consider a situation where the center of mass is all the time located on the y axis, i.e., in $x = 0$. Zero potential energy is chosen in vertical position $y = 0$.



a) (7%) Show that the Lagrangian of the system is

$$L(\theta, \dot{\theta}) = T - V = \frac{m\ell^2\dot{\theta}^2}{4}(1 + \sin^2 \theta) + mg\ell \cos \theta.$$

Hint: $a \sin^2 x + b \cos^2 x = b + (a - b) \sin^2 x$.

b) (7%) Find the equation of motion (i.e., the Lagrange equation).

c) (5%) If the oscillation amplitude is small, the system is a harmonic oscillator. Show this by including only linear terms in the equation of motion, and determine the oscillation frequency ω .

d) (6%) Find the Hamiltonian $H = p_\theta \dot{\theta} - L$ expressed as a function of the canonical variables θ and p_θ .

4 (5%) A point particle moving in two dimensions collides with a hard disk with radius a . If the impact parameter is $s = a/\sqrt{2}$, what is the scattering angle θ ?

The general situation is illustrated in the figure below. The scattering angle θ is defined as the angle between the point particle's incoming and outgoing direction.

